



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07D 493/04, A61K 31/335, C07D 405/06, C07D 405/14, C07D 413/06, C07D 417/06	A1	(11) International Publication Number: WO 00/66589 (43) International Publication Date: 09 November 2000 (09.11.2000)
------------------------------------------------------------------------------------------------------------------------------------------	-----------	---------------------------------------------------------------------------------------------------------------------------------------

(21) International Application Number: PCT/IB00/00657

(22) International Filing Date: 01 May 2000 (01.05.2000)

(30) Priority Data:

100 15 836.6 27 March 2000 (27.03.2000) DE
199 21 086.1 30 April 1999 (30.04.1999) DE
199 54 228.7 04 November 1999 (04.11.1999) DE

(60) Parent Application or Grant

SCHERING AKTIENGESELLSCHAFT [/]; (). KLAR, Ulrich
[/]; (). SCHWEDE, Wolfgang [/]; (). SKUBALLA, Werner [/];
(). BUCHMANN, Bernd [/]; (). HOFFMANN, Jens [/];
(). LICHTNER, Rosemarie [/]; (). KLAR, Ulrich [/];
(). SCHWEDE, Wolfgang [/]; (). SKUBALLA, Werner [/];
(). BUCHMANN, Bernd [/]; (). HOFFMANN, Jens [/];
(). LICHTNER, Rosemarie [/]; (). SOPP, John, A. ; ().

Published

(54) Title: 6-ALKENYL-, 6-ALKINYL- AND 6-EPOXY-EPOTHILONE DERIVATIVES, PROCESS FOR THEIR PRODUCTION, AND THEIR USE IN PHARMACEUTICAL PREPARATIONS

(54) Titre: DERIVES 6-ALCENYL-, 6-ALKINYL-, ET 6-EPOXY-EPOTHILONE, PROCEDE DE PREPARATION ET LEUR UTILISATION DANS DES PREPARATIONS PHARMACEUTIQUES

(57) Abstract

This invention describes the new 6-alkenyl- and 6-alkinyl- epothilone derivatives of general formula (I) in which R1a_z, R1b_z, R2a_z, R3a_z, R3b_z, R4_z, R5_z, R6_z, R7_z, A, Y, D, E, G, Y and Z have the meanings that are indicated in the description. The new compounds interact with tubulin by stabilizing microtubuli that are formed. They are able to influence the cell-splitting in a phase-specific manner and thus find use in treating diseases or conditions associated with the need for cell growth, division and/or proliferation. Thus the compounds are suitable for treating malignant tumors, for example, ovarian, stomach, colon, adeno-, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia. In addition, they are suitable for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases (such as psoriasis, arthritis). Methods of use and preparation of the compounds are also described.

(57) Abrégé

L'invention concerne de nouveaux dérivés 6-alcényl- et 6-alkinyl-épothilone de formule générale (I), dans laquelle R1a_z, R1b_z, R2a_z, R3a_z, R3b_z, R4_z, R5_z, R6_z, R7_z, A, Y, D, E, G, Y et Z sont tels que définis dans le descriptif. Les nouveaux composés interagissent avec la tubuline pour stabiliser les microtubules qui sont formées. Ils peuvent avoir une influence sur la division cellulaire d'une manière spécifique à une phase et trouvent ainsi une application dans le traitement de maladies ou de conditions associées au besoin de croissance, de division et/ou de prolifération cellulaire. Les composés sont donc appropriés dans le traitement de tumeurs malignes, telles que des carcinomes de l'ovaire, de l'estomac, du côlon, du sein, du poumon, de la tête, du cou ou des adéno-carcinomes, des mélanomes malins, une leucémie lymphocytaire ou myélocytaire aiguë. Les composés sont en outre appropriés dans la thérapie anti-angiogénèse ainsi que dans le traitement de maladies inflammatoires chroniques (telles que le psoriasis ou l'arthrite). L'invention concerne également des méthodes d'utilisation et de préparation des composés.

ATTORNEY DOCKET NUMBER: 5624-262-999
SERIAL NUMBER: 09/280,210
REFERENCE: CBJ

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 493/04, 405/06, 417/06, 405/14, 413/06, A61K 31/335		A1	(11) International Publication Number: WO 00/66589
			(43) International Publication Date: 9 November 2000 (09.11.00)
(21) International Application Number: PCT/IB00/00657		Wolfgang [DE/DE]; Klosterheider Weg 35, D-13467 Berlin (DE). SKUBALLA, Werner [DE/DE]; Mattersburger Weg 12, D-13465 Berlin (DE). BUCHMANN, Bernd [DE/DE]; Erdmannstrasse 44, D-16540 Hohen Neuendorf (DE). HOFFMANN, Jens [DE/DE]; Grossstückenfeld 27, D-16567 Mühlenbeck (DE). LICHTNER, Rosemarie [DE/DE]; Belziger Strasse 39, D-10823 Berlin (DE).	
(22) International Filing Date: 1 May 2000 (01.05.00)			
(30) Priority Data: 199 21 086.1 30 April 1999 (30.04.99) DE 199 54 228.7 4 November 1999 (04.11.99) DE 100 15 836.6 27 March 2000 (27.03.00) DE			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 199 21 086.1 (CON) Filed on 30 April 1999 (30.04.99) US 199 54 228.7 (CON) Filed on 4 November 1999 (04.11.99) US 100 15 836.6 (CON) Filed on 27 March 2000 (27.03.00)			
(71) Applicant (for all designated States except US): SCHERING AKTIENGESELLSCHAFT [DE/US]; D-13342 Berlin (DE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): KLAR, Ulrich [DE/DE]; Isegrimsteig 8a, D-13503 Berlin (DE). SCHWEDE,		(74) Agents: SOPP, John, A. et al.; Millen, White, Zelano & Branigan, P.C., Suite 1400, Arlington Courthouse Plaza 1, 2200 Clarendon Boulevard, Arlington, VA 22201 (US).	
		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: 6-ALKENYL-, 6-ALKINYL- AND 6-EPOXY-EPOTHILONE DERIVATIVES, PROCESS FOR THEIR PRODUCTION, AND THEIR USE IN PHARMACEUTICAL PREPARATIONS			
<div style="text-align: center;"><p>(I)</p></div>			
(57) Abstract			
<p>This invention describes the new 6-alkenyl- and 6-alkinyl- epothilone derivatives of general formula (I) in which R^{1a}, R^{1b}, R^{2a}, R^{3a}, R^{3b}, R⁴, R⁵, R⁶, R⁷, A, Y, D, E, G, Y and Z have the meanings that are indicated in the description. The new compounds interact with tubulin by stabilizing microtubuli that are formed. They are able to influence the cell-splitting in a phase-specific manner and thus find use in treating diseases or conditions associated with the need for cell growth, division and/or proliferation. Thus the compounds are suitable for treating malignant tumors, for example, ovarian, stomach, colon, adeno-, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia. In addition, they are suitable for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases (such as psoriasis, arthritis). Methods of use and preparation of the compounds are also described.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

1

Background of the Invention

in, e.g., Angew. Chem. [Applied Chem.], 1996, 108, 1671-1673.

Because of their in-vitro selectivity for breast cell lines and intestinal cell lines and their significantly higher activity against P-glycoprotein-forming multiresistant tumor lines in comparison to taxol as well as their physical properties that are superior to those of taxol, e.g., a water solubility that is higher by a factor of 30, this novel structural class is especially advantageous for the development of a pharmaceutical agent for treating malignant tumors.

5

10

15

20

25

30

35

40

45

50

55

The natural substances are not sufficiently stable either chemically or metabolically for the development of pharmaceutical agents. To eliminate these drawbacks, modifications to the natural substance are necessary. Such modifications are possible only with a total-synthesis approach and require synthesis strategies that make possible a broad modification of the natural substance. The purpose of the structural changes is also to increase the therapeutic range. This can be done by improving the selectivity of the action and/or increasing the active strength and/or reducing undesirable toxic side-effects, as they are described in Proc. Natl. Acad. Sci. USA 1998, 95, 9642-9647.

The total synthesis of epothilone A is described by Schinzer et al. in Chem. Eur. J. 1996, 2, No. 11, 1477-1482 and in Angew. Chem. 1997, 109, No. 5, pp. 543-544). Epothilone derivatives were already described by Höfle et al. in WO 97/19086. These derivatives were produced starting from natural epothilone A or B. Also, epothilones C and D (double bond between carbon atoms 12 and 13: epothilone C = deoxyepothilone A; epothilone D = deoxyepothilone B) are described as possible starting products for this purpose.

Another synthesis of epothilone and epothilone derivatives was described by Nicolaou et al. in Angew. Chem. 1997, 109, No. 1/2, pp. 170-172. The synthesis of epothilone A and B and

several epothilone analogues was described in Nature, Vol. 387,
1997, pp. 268-272; and the synthesis of epothilone A and its
derivatives was described in J. Am. Chem. Soc., Vol. 119, No. 34,
1997, pp. 7960-7973 as well as the synthesis of epothilone A and
B and several epothilone analogues in J. Am. Chem. Soc., Vol.
119, No. 34, 1997, pp. 7974-7991 also by Nicolaou et al.

Nicolaou et al. also describe in Angew. Chem. 1997, 109, No.
19, pp. 2181-2187 the production of epothilone A analogues using
combinative solid-phase synthesis. Several epothilone B
analogues are also described there.

Epothilone derivatives, in some cases also epothilone C and
D, are also described in patent applications WO 99/07692, WO
99/02514, WO 99/01124, WO 99/67252, WO 98/25929, WO 97/19086, WO
98/38192, WO 99/22461 and WO 99/58534.

In the epothilone derivatives previously known, no alkenyl,
alkinyl or epoxy radical was provided on carbon atom 6 (see the
above formula) of the epothilone skeleton.


Summary of the Invention

An object of this invention was to make available new
epothilone derivatives, which are both chemically and
metabolically stable enough for the development of pharmaceutical
agents and which are superior to natural derivatives in terms of

their therapeutic range, their selectivity of action and/or
undesirable toxic side-effects and/or their active strength.

Upon further study of the specification and appended claims,
further objects and advantages of this invention will become
apparent to those skilled in the art.

Included in the invention are new epothilone derivatives and
compounds of general formula I,




I,

in which

R^{1a} , R^{1b} are the same or different and mean hydrogen, C_1 - C_{10}
alkyl, C_6 - C_{12} aryl or C_7 - C_{20} aralkyl, all optionally
substituted, or together a $-(CH_2)_m-$ group with $m = 1,$
2, 3, 4 or 5 or a $-(CH_2)-O-(CH_2)-$ group,

R^{2a} means hydrogen, C_1 - C_{10} alkyl, C_6 - C_{12} aryl or C_7 - C_{20}
aralkyl, all optionally substituted,

$-(CH_2)_{ra}-C\equiv C-(CH_2)_{pa}-R^{26a}$, $-(CH_2)_{ra}-C=C-(CH_2)_{pa}-R^{26a}$,




5

10

R^{2b} means $-(CH_2)_{rb}-C\equiv C-(CH_2)_{pb}-R^{26b}$, $-(CH_2)_{rb}-C=C-(CH_2)_{pb}-R^{26b}$,

15



n means 0 to 5,

20

ra , rb are the same or different and mean 0 to 4,

pa , pb are the same or different and mean 0 to 3,

R^{3a} means hydrogen, C_1 - C_{10} alkyl, C_6 - C_{12} aryl, or C_7 - C_{20}

25

aralkyl, all optionally substituted

R^{14} means hydrogen, OR^{14a} , Hal,

R^{3b} means OH or OPG^{14} ,

30

R^4 means hydrogen, C_1 - C_{10} alkyl, C_6 - C_{12} aryl, or C_7 - C_{20}

aralkyl, all optionally substituted Hal, OR^{25} , CN,

R^{25} means hydrogen, a protective group PG^5 ,

35

R^{26a} , R^{26b} are the same or different and mean hydrogen, C_1 - C_{10}

alkyl, C_6 - C_{12} aryl, or C_7 - C_{20} aralkyl, all optionally

40

substituted, C_1 - C_{10} acyl, or, if pa or $pb > 0$,

additionally a group OR^{27} ,

R^{27} means hydrogen, a protective group PG^6 ,

45

R^5 means hydrogen, C_1 - C_{10} alkyl, C_6 - C_{12} aryl, C_7 - C_{20} aralkyl,

all optionally substituted $(CH_2)_s-T$,

whereby

50

55

5

10

15

20

25

30

35

40

45

50

55

s stands for 1, 2, 3 or 4,

T stands for OR^{22} or Hal,

R^{22} stands for hydrogen or a protective group

PG^4 ,

R^6 , R^7 each mean a hydrogen atom, or taken together an additional bond or an oxygen atom,

G means a group $X=CR^8-$, a bi- or tricyclic aryl radical,

R^8 means hydrogen, halogen, CN, C_1-C_{20} alkyl, C_6-C_{12} aryl, C_7-C_{20} aralkyl, which can all be substituted,

X means an oxygen atom, two alkoxy groups OR^{23} , a C_2-C_{10} -alkylene- α,ω -dioxy group, which can be straight-chain or branched, H/OR^9 or a grouping $CR^{10}R^{11}$,

whereby

R^{23} stands for a C_1-C_{20} alkyl radical, optionally substituted

R^9 stands for hydrogen or a protective group PG^x ,

R^{10}, R^{11} are the same or different and stand for

hydrogen, a C_1-C_{20} alkyl, C_6-C_{12} aryl, or C_7-C_{20} aralkyl radical, all optionally substituted, or R^{10} and R^{11} together with the methylene carbon atom jointly stand for a 5- to 7-membered carbocyclic ring,

D-E means a group $-CH_2-CH_2-$, $-O-CH_2-$,

5

10

A-Y means a group O-C(=O), O-CH₂, CH₂C(=O), NR²⁹-C(=O),
NR²⁹-SO₂,

15

R²⁹ means hydrogen, C₁-C₁₀ alkyl,

Z means an oxygen atom or H/OR¹²,

whereby

R¹² is hydrogen or a protective group PG²,

20

Hal means halogen, preferably fluorine, chlorine, bromine
or iodine.


25

The production of the new epothilone derivatives and
compounds can be performed by linkage of three partial fragments
A_f, B_f and C_f which process is a further aspect of the invention.
The interfaces between the fragments are as indicated by the
three lines crossing bands in general formula I'.

30

35

40




I',

45

50

55

A means a C₁-C₆ fragment (epothilone numbering system) of
general formula A-1 or A-2



in which

R^{1a'}, R^{1b'}, R^{2a'} and R^{2b'} have the meanings already mentioned

for R^{1a}, R^{1b}, R^{2a} and R^{2b}, and

R¹³ means CH₂OR^{13a}, CH₂-Hal, CHO, CO₂R^{13b}, COHal,

R^{14'} means hydrogen, OR^{14a}, Hal, OSO₂R^{14b},

R^{13a}, R^{14a} mean hydrogen, SO₂-alkyl, SO₂-aryl, SO₂-aralkyl or
together a -(CH₂)_o- group or together a CR^{15a}R^{15b} group,

R^{13b}, R^{14b} mean hydrogen, C₁-C₂₀ alkyl, C₆-C₁₂ aryl, C₇-C₂₀

aralkyl, each optionally substituted

R^{15a}, R^{15b} are the same or different and mean hydrogen, C₁-C₁₀

alkyl, aryl, C₇-C₂₀ aralkyl or together a -(CH₂)_q group,

o means 2 to 4,

q means 3 to 6,

R³⁰ means hydrogen,

R³¹ means hydroxyl, or

R^{30} , R^{31} together mean an oxygen atom or a C_2 - C_{10} alkylene-


α,ω -dioxy group, which can be straight-chain or branched, or

R^{30} , R^{31} independently mean a C_1 - C_{10} alkoxy group,

including all stereoisomers as well as their mixtures, and

free hydroxyl groups in R^{13} , R^{14} and R^{31} can be etherified or esterified, free carbonyl groups can be ketalized in A-1 or A-2 and R^{13} , converted into an enol ether or reduced, and free acid groups in A-1 or A-2 can be converted into their salts with bases.

B_f stands for a C_7 - C_{12} fragment (epothilone numbering system) of general formula



B_f

in which

D , E , $R^{3a'}$, $R^{4'}$ and $R^{5'}$ have the meanings already mentioned for D , E , R^{3a} , R^4 , and R^5 , and

5

10

10

NOT FURNISHED UPON FILING

15

20

25

30

35

40

45

50

55

5

$R^{7'}$ means a hydrogen atom,

10

R^{20} means halogen, N_3 , NHR^{29} , a hydroxy group, a protected hydroxy group $O-PG^2$, a protected amino group $NR^{29}PG^2$, a C_1-C_{10} alkylsulfonyloxy group, which optionally can be perfluorinated, a benzyloxy group that is optionally substituted by C_1-C_4 alkyl, nitro, chlorine or bromine, an $NR^{29}SO_2CH_3$ group, an $NR^{29}C(=O)CH_3$ group, a $CH_2-C(=O)-CH_3$ group,

15

20

R^{21} means a hydroxy group, halogen, a protected hydroxy group OPG^3 , a phosphonium halide radical $PPh_3^+Hal^-$ ($Ph =$ phenyl; $Hal = F, Cl, Br, I$), a phosphonate radical $P(O)(OQ)_2$ ($Q = C_1-C_{10}$ alkyl or phenyl) or a phosphine oxide radical $P(O)Ph_2$ ($Ph =$ phenyl),

25

30

X means an oxygen atom, two alkoxy groups OR^{23} , a C_2-C_{10} alkylene- α,ω -dioxy group, which can be straight-chain or branched, H/OR^9 or a grouping $CR^{10}R^{11}$,
whereby

35

40

R^{23} stands for a C_1-C_{20} alkyl radical,

R^9 stands for hydrogen or a protective group PG^3 ,

R^{10} , R^{11} are the same or different and stand for

45

hydrogen, a C_1-C_{20} alkyl, C_6-C_{12} aryl, or C_7-C_{20}

aralkyl radical, each optionally substituted, or

R^{10} and R^{11} together with the methylene carbon atom

50

55

5

10

commonly stand for a 5- to 7-membered carbocyclic ring.

15

20

As alkyl groups R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^{3a} , R^4 , R^5 , R^8 , R^{10} , R^{11} , R^{13a} , R^{13b} , R^{14a} , R^{14b} , R^{15a} , R^{15b} , R^{17} , R^{19} , R^{23} , R^{26a} , R^{26b} , R^{28a} , R^{28b} and R^{29} , straight-chain or branched-chain alkyl groups with 1-20 carbon atoms can be considered, such as, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl, heptyl, hexyl, and decyl.

25

30

Alkyl groups R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^{3a} , R^4 , R^5 , R^8 , R^{10} , R^{11} , R^{13a} , R^{13b} , R^{14a} , R^{14b} , R^{15a} , R^{15b} , R^{17} , R^{19} , R^{23} , R^{26a} , R^{26b} , R^{28a} and R^{28b} can be perfluorinated or substituted by 1-5 halogen atoms, hydroxy groups, C_1 - C_4 alkoxy groups, C_6 - C_{12} aryl groups (which can be substituted by 1-3 halogen atoms).

35

40

45

50

As aryl radicals above and below, particularly R^{1a} , R^{1b} , R^{2a} , R^{2b} , R^{3a} , R^4 , R^5 , R^8 (R^{10} , R^{11} , R^{13a} , R^{13b} , R^{14a} , R^{14b} , R^{15a} , R^{15b} , R^{26a} , R^{26b} , R^{28a} and R^{28b} , substituted and unsubstituted carbocyclic or heterocyclic radicals with one or more heteroatoms, such as, e.g., phenyl, naphthyl, furyl, thienyl, pyridyl, pyrazolyl, pyrimidinyl, oxazolyl, pyridazinyl, pyrazinyl, quinolyl, thiazolyl, benzothiazolyl and benzoxazolyl, which can be substituted in one or more places by halogen, OH, O-alkyl, CO_2H , CO_2 -alkyl, ($-NH_2$, $-NO_2$, $-N_3$, $-CN$, C_1 - C_{20} alkyl, C_1 - C_{20} acyl, C_1 - C_{20} acyloxy groups, are suitable examples.

55

5

10

15

20

25

As bi- and tricyclic aryl radicals G or G', substituted and unsubstituted, carbocyclic or heterocyclic radicals with one or more heteroatoms, such as, e.g., naphthyl, anthryl, benzothiazolyl, benzoxazolyl, benzimidazolyl, quinolyl, isoquinolyl, benzoxazinyl, benzofuranyl, indolyl, indazolyl, quinoxalinyl, tetrahydroisoquinolyl, tetrahydroquinolyl, thienopyridinyl, pyridopyridinyl, benzopyrazolyl, benzotriazolyl, dihydroindolyl, which can be substituted in one or more places by halogen, OH, O-alkyl, CO₂H, CO₂-alkyl, -NH₂, -NO₂, -N₃, -CN, C₁-C₂₀ alkyl, C₁-C₂₀ acyl, C₁-C₂₀ acyloxy groups, are suitable.

30

35

40

The aralkyl groups in R^{1a}, R^{1b}, R^{2a}, R^{2b}, R^{3a}, R⁴, R⁵, R⁸, R¹⁰, R¹¹, R^{13a}, R^{13b}, R^{14a}, R^{14b}, R^{15a}, R^{15b}, R^{26a}, R^{26b}, R^{28a} and R^{28b} can contain in the ring(s) up to 14 C atoms, preferably 6 to 10, and in the alkyl chain 1 to 8, preferably 1 to 4 atoms. As aralkyl radicals, for example, benzyl, phenylethyl, naphthylmethyl, naphthylethyl, furylmethyl, thienylethyl and pyridinylpropyl are suitable. The rings can be substituted in one or more places by halogen, OH, O-alkyl, CO₂H, CO₂-alkyl, -NO₂, -N₃, -CN, C₁-C₂₀ alkyl, C₁-C₂₀ acyl, C₁-C₂₀ acyloxy groups.

45

50

The alkoxy groups that are contained in R³⁰, R³¹ and X in general formula I are in each case to contain 1 to 20 carbon atoms, whereby methoxy, ethoxy, propoxy, isopropoxy and t-butyloxy groups are preferred.

55

5

10

15

20

As representatives of all the protective groups PG (i.e., each of the PG groups including those with a superscript), alkyl- and/or aryl-substituted silyl, C₁-C₂₀ alkyl, C₄-C₇ cycloalkyl, which in addition can contain an oxygen atom in the ring, aryl, C₇-C₂₀ aralkyl, C₁-C₂₀ acyl, aroyl and C₁-C₂₀ alkoxy carbonyl can be mentioned. Other protective groups are known in the art and can be used as known.

25

30

35

40

As alkyl, silyl and acyl radicals for protective groups PG, the radicals that are known to one skilled in the art are suitable. Preferred are alkyl or silyl radicals that can be easily cleaved from the corresponding alkyl and silyl ethers, such as, for example, the methoxymethyl, methoxyethyl, ethoxyethyl, tetrahydropyranyl, tetrahydrofuranyl, trimethylsilyl, triethylsilyl, tert-butyldimethylsilyl, tert-butyldiphenylsilyl, tribenzylsilyl, triisopropylsilyl, benzyl, para-nitrobenzyl, para-methoxybenzyl radical as well as alkylsulfonyl and arylsulfonyl radicals. As acyl radicals, e.g., formyl, acetyl, propionyl, isopropionyl, pivalyl, butyryl or benzoyl, which can be substituted with amino and/or hydroxy groups, are suitable.

45

50

55

As amino protective groups, the radicals that are known to one skilled in the art are suitable. For example, the Alloc-,

Boc-, Z-, benzyl, f-Moc, Troc, Stabase or Benzostabase group can be mentioned.

Acyl groups PG can contain 1 to 20 carbon atoms, whereby formyl, acetyl, propionyl, isopropionyl and pivalyl groups are preferred.

Index m in the alkylene group that is formed from R^{1a} and R^{1b} preferably stands for 1, 2, 3 or 4.

The C₂-C₁₀ alkylene- α,ω -dioxy group that is possible for R³⁰, R³¹, U, V, W and X is preferably an ethyleneketal or neopentylketal group.

The compounds that are mentioned below are preferred according to the invention:

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-Dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R), 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-
pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-
en-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-
dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-
pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-
en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-
4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-
pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-
en-1-yl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-
4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2-
(2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-
oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-
oxacyclopropyl-1-ethyl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -

50

55

5

10

8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -
cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-
4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -
cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-
dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -
cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-
4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-Dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -
cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-Dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-
pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
en-1-yl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-
4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2-
(2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-
(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-
(oxacyclopropylmethyl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -
8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-
dione

30

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-
pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-
(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-
(oxacyclopropylmethyl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2-
(2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-
(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-
(oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) -
8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-
dione

15

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2-
(2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7-
(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-
(oxacyclopropylmethyl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) -
8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-
dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-
methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-
in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-
in-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-
methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-
in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-
in-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

45

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, (10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-

45

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
in-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-
methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-
en-1-yl) -cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
en-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-
methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-
en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
en-1-yl) -3- (2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-
methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-
en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

30

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(but-3-in-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

45

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-

35

tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5, 5, 9, 13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

30

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

55

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-
(oxacyclopropylmethyl) - 3 - (1-fluoro-2 - (2-methyloxazol-4-
yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-
dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-chloro-2-
(2-methyloxazol-4-yl)ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7-
(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-
(oxacyclopropylmethyl) - 3 - (1-chloro-2 - (2-methyloxazol-4-
yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-
dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-methyl-2 - (2-
pyridyl)ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7 - (but-3-
in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10 - (but-3-
in-1-yl) - 3 - (1-methyl-2 - (2-pyridyl)ethenyl) - 8, 8-trimethylene-
12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-fluoro-2 - (2-
pyridyl)ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7 - (prop-2-
in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10 - (prop-2-
in-1-yl) - 3 - (1-fluoro-2 - (2-pyridyl)ethenyl) - 8, 8-trimethylene-
12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione .

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-
1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-
2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-
en-1-yl)-3-(1-methyl-2-(quinolin-2-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-
1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-
2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-
in-1-yl)-3-(1-methyl-2-(quinolin-2-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-
1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-
cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-
methyl-but-2-en-1-yl)-3-(1-methyl-2-(quinolin-2-yl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-
benzoxazol-5-yl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-
cyclohexadec-13-ene-2,6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (quinolin-2-yl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (quinolin-2-yl) -
1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-
ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
in-1-yl) -3- (1-methyl-2- (quinolin-2-yl) -8, 8, 12, 16-tetramethyl-4-
aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (quinolin-2-yl) -
1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-
2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-
en-1-yl) -3- (1-methyl-2- (quinolin-2-yl) -8, 8, 12, 16-tetramethyl-4-
aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (quinolin-2-yl) -
1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-
2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-
in-1-yl) -3- (1-methyl-2- (quinolin-2-yl) -8, 8, 12, 16-tetramethyl-4-
aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (quinolin-2-yl) -
1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

. (4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.10]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.10]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-

50

55

5

10

8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl-3-(2-(2-pyridyl)ethenyl))-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-

35

8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

50

55

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-

35

tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) - ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

55

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-chloro-2- (2-methylthiazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methylthiazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

20

25

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-

45

oxabicyclo[14.1.0]heptadecane-5,9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-

35

tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-

30

oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

45

oxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-

50

55

5

10

15

20

25

30

35

40

45

50

55

trimethylene-12,16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5-trimethylene-9,13-dimethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8-trimethylene-12,16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5-trimethylene-9,13-dimethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8-

trimethylene-12,16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5-trimethylene-9,13-dimethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8-trimethylene-12,16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

5

10

15

20

25

30

35

40

45

50

55

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5, 5-trimethylene-9, 13-dimethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5-trimethylene-9, 13-dimethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-in-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5-trimethylene-9, 13-dimethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5, 5-trimethylene-9, 13-dimethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2, 6-dione

5

10

15

20

25

30

35

40

45

50

55

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12-16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-chloro-2- (2-methyloxazol-4-

5

10

15

20

25

30

35

40

45

50

55

yl)ethenyl)-8,8-trimethylene-12,16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxobicyclo[14.1.0]heptadecane-5,9-dione

20

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-methyl-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

20

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

45

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

45

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

50

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-

55

5

10

15

20

25

30

35

40

45

50

55

yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-

20

oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

40

45


50

55

Production of Partial Fragments A_f:

The partial fragments (synthesis components) of general formula A-1 can be produced starting from the precursors described in WO 99/07692, such as, for example, A-I. This is further explained by way of example in Diagram 1.

Diagram 1

Step a (A-I \Rightarrow A-II):

The hydroxyl group that is protected by PG¹ in A-1 is released. As protective group PG¹, the protective groups

described above and that are known to one skilled in the art,
such as, e.g., the methoxymethyl, methoxyethyl, ethoxyethyl,
tetrahydropyranyl, tetrahydrofuranyl, trimethylsilyl,
triethylsilyl, tert-butyldimethylsilyl, tert-butyldiphenylsilyl,
tribenzylsilyl, triisopropylsilyl, benzyl, para-nitrobenzyl,
para-methoxybenzyl, formyl, acetyl, propionyl, isopropionyl,
pivalyl, butyryl or benzoyl radicals, are suitable. A survey is
found in, e.g., "Protective Groups in Organic Synthesis,"
Theodora W. Green, John Wiley and Sons).

Preferred are those protective groups that can be cleaved
under the action of fluoride, such as, e.g., the trimethylsilyl,
tert-butyldimethylsilyl, tert-butyldiphenylsilyl, tribenzylsilyl,
or triisopropylsilyl radical.

Especially preferred are the tert-butyldimethylsilyl
radical, the triisopropylsilyl radical, and the tert-
butyldiphenylsilyl radical.

As protective groups PG^{8a} and PG^{8b} , the groups that are
already mentioned for PG^7 and together a $-CR^{28a}R^{28b}$ group, in which
 R^{28a} and R^{28b} can be the same or different and mean hydrogen, C_1 - C_{10}
alkyl, aryl, C_7 - C_{20} aralkyl, are suitable.

Preferred are those $-CR^{28a}R^{28b}$ protective groups, in which R^{28a}
and R^{28b} mean C_1 - C_8 alkyl, or R^{28a} means hydrogen and R^{28b} means
aryl.

Especially preferred is a $-C(CH_3)_2$ group.

Protective group PG' is cleaved according to a process that is known to one skilled in the art. This is a silyl ether, thus suitable for the cleavage is the reaction with fluorides, such as, for example, tetrabutylammonium fluoride, hydrogen fluoride-pyridine complex, potassium fluoride or the use of dilute mineral acids, the use of catalytic amounts of acids, such as, e.g., para-toluenesulfonic acid, para-toluenesulfonic acid-pyridinium salt, camphorsulfonic acid in alcoholic solutions, preferably in ethanol or isopropanol.

Step b (A-II \rightarrow A-III):

The oxidation of the primary alcohol in A-II to aldehyde A-III is carried out according to the methods that are known to one skilled in the art. For example, oxidation with pyridinium chlorochromate, pyridinium dichromate, chromium trioxide-pyridine complex, oxidation according to Swern or related methods, e.g., with use of oxalyl chloride in dimethyl sulfoxide, the use of Dess-Martin periodinane, the use of nitrogen oxides, such as, e.g., N-methyl-morpholino-N-oxide in the presence of suitable catalysts, such as, e.g., tetrapropylammonium perruthenate in inert solvents, can be mentioned. Preferred is the oxidation

5
10 according to Swern, as well as with N-methyl-morpholino-N-oxide
using tetrapropylammonium perruthenate.

15 Step c (A-III \rightarrow A-IV):

The reaction of aldehydes A-III to alcohols of formula A-IV
is carried out with organometallic compounds of theoretical
20 formula $M-CH_2R^{2a'}$, in which M stands for indium, an alkali metal,
preferably lithium or a divalent metal MX, in which X represents
a halogen, and radical $R^{2a'}$ has the above-mentioned meanings. As
25 a divalent metal, magnesium and zinc are preferred; as halogen, X
is preferably chlorine, bromine and iodine.

30 Step d (A-IV \rightarrow A-V):

The oxidation of the secondary alcohol in A-IV to ketone A-V
35 is carried out according to the conditions that are mentioned
under step b). Preferred is the oxidation with N-methyl-
morpholino-N-oxide with use of tetrapropylammonium perruthenate.

40 Step e (A-V \rightarrow A-VI):

45 For optional introduction of a radical $R^{2b'}$, which, except
for hydrogen, can have the already mentioned meanings, the ketone
of general formula A-V is converted into the enolate with M in

50

55

5

10

the meaning of the counter-cation with use of strong bases, such as preferably lithium diisopropylamide.

15

Step f (A-VI - A-VII):

20

The enolate of formula A-VI is reacted with a compound of general formula $X-R^{2b'}$, in which X represents a halogen or another leaving group, such as, for example, an alkylsulfonate or arylsulfonate. As halogen, X is preferably chlorine, bromide and iodine.

25

30

The partial fragments (synthesis components) of general formula A-2 can be produced by known methods, such as described in Angew. Chemie 1996, 108, 2976-2978. Another process is shown in Diagram 2:

35


40

45

50

55

Diagram 2



Step a (A-VIII \Rightarrow A-IX).

The oxidation of the primary alcohol in A-VIII to aldehyde A-IX is carried out according to the methods that are described for Diagram 1, step b. The oxidation according to Swern and with N-methyl-morpholino-N-oxide with use of tetrapropylammonium perruthenate is preferred.

Step b (A-IX \Rightarrow A-X):

The carbonyl group in A-IX can optionally be converted into a ketal according to the methods that are known to one skilled in the art.

Step c (A-X \Rightarrow A-XI):

The hydroxyl group that is protected in A-X by PG⁹ is released. As protective group PG⁹, the protective groups that are described under Diagram 1, step a are suitable. Preferred are those protective groups that can be cleaved under the action of fluoride, such as, e.g., the trimethylsilyl, tert-butyldimethylsilyl, tert-butyldiphenylsilyl, tribenzylsilyl, or triisopropylsilyl radical.

Especially preferred is the tert-butyldimethylsilyl radical, the triisopropylsilyl radical and the tert-butyldiphenylsilyl radical.

Step d (A-XI \Rightarrow A-XII):

The oxidation of the primary alcohol in A-IX to aldehyde A-XII is carried out according to the methods that are described for Diagram 1, step b. The oxidation according to Swern and with N-methyl-morpholino-N-oxide with use of tetrapropylammonium perruthenate is preferred.

Step e (A-XII \Rightarrow A-XIII):

The introduction of radicals R^{2a'} and/or R^{2b'} and production of ketone A-XIII are carried out as described under Diagram 1 in steps c to f.

5

Production of Partial Fragments B_f:

10

The partial fragments (synthesis components) of general formula B_f can be produced by known methods, such as described in WO 99/07692.

15

Production of Partial Fragments C_f:

20

The partial fragments (synthesis components) of general formula C_f can be produced as described in DE 197 51 200.3, DE 199 07 480.1 and WO 99/07692 as well as in PCT/EP00/01333 and Example 21.

25

Production of partial fragments A_f, B_f and C_f and their cyclization to I is also carried out analogously to what is described in WO 99/07692 for numerous epothilone derivatives, with the difference that in the known derivatives in 6-position, there is no unsaturated radical. WO 99/07692 already confirms the general usability of the synthesis principle that is described below for the compounds according to the invention. In addition, numerous synthesis components of general formulas A_f, B_f and C_f are described in WO 99/07692, with which, optionally in modified form in the case of the substitution according to the invention at carbon 6, other compounds of general formula I claimed here can be obtained. Synthesis components of general formula C_f, in which as R⁶, a halogen atom, especially a

30

35

40


45

50

55


fluorine, chlorine or bromine atom, is present, are the subject
of DE 199 07 480.1 as well as PCT/EP00/01333.


Partial fragments of general formula AB



in which $R^{1a'}$, $R^{1b'}$, $R^{2a'}$, $R^{2b'}$, R^{3a} , R^4 , R^5 , R^{13} , R^{14} , R^{30} , R^{31} , V and Z
have the already mentioned meanings, and $PG^{14'}$ represents a
hydrogen atom or a protective group PG , can be obtained from the
above-mentioned fragments A_f and B_f according to the process that
is shown in Diagram 3.

Diagram 3






Step aa ($A_f + B_f \rightarrow AB$):

Compound B_f , in which W has the meaning of an oxygen atom and optionally present additional carbonyl groups are protected, is alkylated with the enolate of a carbonyl compound of general formula A_f . The enolate is produced by action of strong bases, such as, e.g., lithium diisopropylamide, lithium hexamethyldisilazane at low temperatures.


Partial Fragments of General Formula BC



BC,


in which R^{3a} , R^4 , R^5 , R^6 , R^7 , R^{20} , D , E , G and W have the already mentioned meanings, are obtained from previously described fragments B_f and C_f according to the process that is shown in Diagram 4.

Diagram 4



Step ab ($B_f + C_f \rightarrow BC$):



Compound C_f , in which R^{21} has the meaning of a Wittig salt, and optionally present additional carbonyl groups are protected, is deprotonated by a suitable base, such as, e.g., *n*-butyllithium, lithium diisopropylamide, potassium *tert*-butanolate, sodium or lithium-hexamethyldisilazide and reacted with a compound B_f , in which V has the meaning of oxygen, and W has the meaning of two alkoxy groups OR^{19} , a C_2 - C_{10} -alkylene- α,ω -dioxy group, which can be straight-chain or branched or has H/OR^{18} .

Partial Fragments of General Formula ABC (AB + C_r)

in which $R^{1a'}$, $R^{1b'}$, $R^{2a'}$, $R^{2b'}$, R^{3a} , R^4 , R^5 , R^6 , R^7 , R^{13} , R^{14} , R^{30} , R^{31} ,

D, E, G and Z have the meanings already mentioned, and $PG^{14'}$


represents a hydrogen atom or a protective group PG, are obtained from the previously described fragments AB and C according to the process that is shown in Diagram 5 and Diagram 6.



Step ac ($AB + C_f = ABC$):

Compound C_f , in which R^{21} has the meaning of a Wittig salt, and optionally present additional carbonyl groups are optionally protected, is deprotonated by a suitable base, such as, e.g., n-butyllithium, lithium diisopropylamide, potassium tert-butanolate, sodium or lithium-hexamethyldisilazide and reacted with a compound AB, in which V has the meaning of an oxygen atom.

Diagram 6



Step ad ($A_f + BC = ABC$):

Compound BC, in which W has the meaning of an oxygen atom and optionally present additional carbonyl groups are protected,

is alkylated with the enolate of a carbonyl compound of general formula A_f . The enolate is produced by action of strong bases, such as, e.g., lithium diisopropylamide, lithium hexamethyldisilazane at low temperatures.

Step ae (ABC-1 \rightarrow I):

Compounds ABC-1, in which R^{13} represents a carboxylic acid CO_2H and R^{20} represents a hydroxyl group or an amino group, are reacted according to the methods that are known to one skilled in the art for the formation of large macrolides or macrolactams to compounds of formula I, in which A-Y has the meaning of an O-(C=O) group or $NR^{29}-C(=O)$ group. For example, preferred for lactone formation is the method that is described in "Reagents for Organic Synthesis, Vol. 16, p. 353" with use of 2,4,6-trichlorobenzoic acid chloride and suitable bases, such as, e.g., triethylamine, 4-dimethylaminopyridine, sodium hydride. For example, preferred for the lactam formation is the reaction of the amino acid (R^{13} a carboxylic acid CO_2H and R^{20} an NHR^{29} group) with diphenylphosphorylazide in the presence of a base.

Step af (ABC-1 \rightarrow I):

Compounds ABC-1, in which R^{13} represents a group CH_2OH and R^{20} represents a hydroxyl group, can be reacted preferably with use

5
10 of triphenylphosphine and azodiester, such as, for example, azodicarboxylic acid diethyl ester, to compounds of formula I, in which A-Y has the meaning of an O-CH₂ group.

15 Compounds ABC, in which R¹³ represents a group CH₂-Hal or CH₂OSO₂- alkyl or CH₂OSO₂- aryl or CH₂OSO₂- aralkyl and R²⁰ represents a hydroxyl group, can be cyclized to compounds of
20 formula I, in which A-Y has the meaning of an O-CH₂ group, after deprotonation with suitable bases, such as, for example, sodium hydride, n-butyllithium, 4-dimethylaminopyridine, Hünig base,
25 alkylhexamethyldisilazanes.

30 Step ag (ABC-2 → 1):

35 Compounds ABC-2, in which R³⁰ and R³¹ together represent an oxygen atom and R²⁰ represents an NR²⁹SO₂CH₃ group, can be cyclized at low temperatures to sulfonamide I, in which A-Y has the
40 meaning of an NR²⁹SO₂ group, by action of strong bases, such as, e.g., lithium diisopropylamide, lithium hexamethyldisilazane.

45 Step ah (ABC-2 → 1):

50 Compounds ABC-2, in which R³⁰ and R³¹ together represent an oxygen atom and R²⁰ represents an O-C(=O)CH₃ group, can be
55 cyclized at low temperatures to lactone I, in which A-Y has the

5
10 meaning of an O-C(=O) group, by action of strong bases, such as,
e.g., lithium diisopropylamide, or alkali hexamethyldisilazane.

15 Step ah (ABC-2 \rightarrow I):

Compounds ABC-2, in which R³⁰ and R³¹ together represent an
oxygen atom and R²⁰ represents a CH₂C(=O)CH₃ group, can be
20 cyclized at low temperatures to lactone I, in which A-Y has the
meaning of a CH₂C(=O) group by action of strong bases such as,
e.g., lithium diisopropylamide, alkali hexamethyldisilazane.
25

Introduction of the Nitrogen Group for R²⁰:

30 Amino group NHR²⁹ can be introduced in the stage of the C_r-
fragment, the BC-fragment or the ABC-fragment according to the
methods that are known to one skilled in the art. Preferred is
35 the production from the azide (R²⁰ = N₃), which is first converted
into the phosphaimine according to the methods that are known to
one skilled in the art, preferably with use of a phosphine such
40 as, for example, triphenylphosphine; the phosphaimine is then
converted in the presence of water into the amine (R²⁰ = NHR²⁹)
that is optionally to be protected for the further course of
45 reaction. The introduction of the azide can be carried out with
use of the Mitsunobu reaction in the presence of metal azides,
preferably sodium or tin azide or by substitution of a suitable
50

5

10

15

leaving group, such as, for example, a chlorine, bromine or
iodine atom, an alkylsulfonyloxy group, a perfluorinated
alkylsulfonyloxy group, an arylsulfonyloxy group or an
aralkylsulfonyloxy group by azides.

20

The flexible functionalization of described components A_r,
B_r and C_r also ensures a linkage sequence that deviates from the
above-described process and that leads to components ABC. These
processes are listed in the following table:

25

30

35

40

45

50


55

Possible Linkages	Linkage Methods a to e	Prerequisites
$A_f + B_f \Rightarrow A_f - B_f$	a: Aldol (see Diagram 3)	Z = W = oxygen
$B_f + C_f \Rightarrow B_f - C_f$	b: Wittig (analogously to Diagram 4) e: McMurry	U = oxygen and R^{21} = Wittig salt, phosphine oxide or phosphonate U = V = oxygen
$A_f + C_f \Rightarrow A_f - C_f$	c: Esterification (e.g., 2,4,6-trichlorobenzoyl chloride and 4-dimethylamino-pyridine) d: Etherification (e.g., according to Mitsunobu) f. Amide formation (e.g., with $(PhO)_2P(O)N_3$) in the presence of a base in an inert solvent g. Ketone formation by aldol reaction with a strong base. h. Sulfonamide formation in the presence of a strong base. i. Amide formation in the presence of a strong base.	$R^{13} = CO_2R^{13b}$ or COHal and $R^{20} = \text{hydroxyl}$ $R^{13} = CH_2OH$ and $R^{20} = \text{hydroxyl or } OSO_2\text{-alkyl or } OSO_2\text{-aryl or } OSO_2\text{-aralkyl}$ $R^{13} = CO_2R^{13b}$ or COHal and $R^{20} = NHR^{29}$ $R^{20} = CH_2C(=O)CH_3$ and $R^{30}, R^{31} = \text{oxygen}$ $R^{20} = NR^{29}SO_2CH_3$ and $R^{30}, R^{31} = \text{oxygen}$ $R^{20} = NR^{29}C(=O)CH_3$ and $R^{30}, R^{31} = \text{oxygen}$

According to these processes, components A_f , B_f and C_f can be

linked as indicated in Diagram 7:

Diagram 7



Free hydroxyl groups in I, A_f , B_f , C_f , AB, BC, ABC can be further functionally modified by etherification or esterification, free carbonyl groups by ketalization, enol ether formation or reduction.

The invention relates to all stereoisomers of these compounds and also mixtures thereof.

5

10

The invention also relates to all prodrug formulations of these compounds, i.e., all compounds that release *in vivo* a bioactive active ingredient component of general formula I.

15

Biological Actions and Applications of the New Derivatives:

20

25

30

35

40

45

50

55

The new compounds of formula I are valuable pharmaceutical agents. They interact with tubulin by stabilizing microtubuli that are formed and are thus able to influence the cell-splitting in a phase-specific manner. Accordingly, the compounds find use in treating diseases or conditions associated with cell growth, division and/or proliferation. They have particular application with quick-growing, neoplastic cells, whose growth is largely unaffected by intercellular regulating mechanisms. Active ingredients having properties of this type, including the compounds of this invention, are suitable for treating malignant tumors. As applications, there can be mentioned, for example, the therapy of ovarian, stomach, colon, adeno-, breast, lung, head and neck carcinomas, malignant melanoma, acute lymphocytic and myelocytic leukemia, particularly against tumors associated therewith. The compounds according to the invention are additionally suitable owing to their properties for anti-angiogenesis therapy as well as for treatment of chronic inflammatory diseases, such as, for example, psoriasis, multiple

5
10 sclerosis or arthritis. The compounds of the invention have
utilities analogous to those for known epothilone derivatives,
discussed above, but with the advantages discussed above and
15 below. As with the known epothilones, the new compounds can be
administered for and in a manner analogous to known drugs such as
Taxol, but modified to take advantage of their particular
20 superior properties thereover.

In other embodiments, to avoid uncontrolled proliferation of
cells and/or for better compatibility of medical implants, the
25 new compounds or pharmaceutical formulations containing them can
be applied or introduced into the polymer materials that are used
for these purposes.
30

The compounds according to the invention can be used alone
or in combination with other principles and classes of substances
35 that can be used in tumor therapy to achieve additive or
synergistic actions. As examples, there can be mentioned the
combination with

- 40
- Platinum complexes, such as, e.g., cis-platinum,
carboplatinum,
 - 45 • intercalating substances, e.g., from the class of
anthracyclines, such as, e.g., doxorubicin or from the
class of anthrapyrazoles, such as, e.g., Cl-941,
- 50

5

103

10

15

20

25

30

35

40

45

50

55

- substances that interact with tubulin, e.g., from the class of vinca-alkaloids, such as, e.g., vincristine, vinblastine or from the class of taxanes, such as, e.g., taxol, taxotere or from the class of macrolides, such as, e.g., rhizoxin or other compounds, such as, e.g., colchicine, combretastatin A-4, discodermolide and its analogs,
- DNA topoisomerase inhibitors, such as, e.g., camptothecin, etoposide, topotecan, teniposide,
- folate- or pyrimidine-antimetabolites, such as, e.g., lometrexol, gemcitabin,
- DNA-alkylating compounds, such as, e.g., adozelesin, dystamycin A,
- inhibitors of growth factors (e.g., of PDGF, EGF, TGF β , EGF), such as, e.g., somatostatin, suramin, bombesin antagonists,
- inhibitors of protein tyrosine kinases or protein kinases A or C, such as, e.g., erbstatin, genistein, staurosporine, ilmofofosine, 8-Cl-cAMP,
- antihormones from the class of antigestagens, such as, e.g., mifepristone, onapristone or from the class of antiestrogens, such as, e.g., tamoxifen or from the

5

104

10

class of antiandrogens, such as, e.g., cyproterone acetate;

15

- metastases-inhibiting compounds, e.g., from the class of eicosanoids, such as, e.g., PGI₂, PGE₁, 6-oxo-PGE₁ as well as their more stable derivatives (e.g., iloprost, cicaprost, misoprostol),

20

- inhibitors of oncogenic RAS proteins, which influence the mitotic signal transduction, such as, for example, inhibitors of the farnesyl-protein-transferase,

25

- natural or synthetically produced antibodies, which are directed against factors or their receptors, which promote tumor growth, such as, for example, the erbB2 antibody.

30

35

The invention also relates to pharmaceutical agents that are based on pharmaceutically compatible compounds, i.e., compounds of general formula I that are nontoxic in the doses used, optionally together with commonly used adjuvants and vehicles.

40

In other embodiments, the compounds according to the invention can be encapsulated with liposomes or included in an α -, β - or γ -cyclodextrin clathrate.

45

50

According to methods of galenicals that are known in the art, the compounds according to the invention can be processed into pharmaceutical preparations for enteral, percutaneous,

55

5

10

15

parenteral or local administration. They can be administered in the form of tablets; coated tablets; gel capsules; granulates; suppositories; implants; injectable, sterile, aqueous or oily solutions; suspensions or emulsions; ointments; creams and gels, for example.

20

25

30

In this case, the active ingredient or ingredients can be mixed with the adjuvants that are commonly used in galenicals, such as, e.g., gum arabic, talc, starch, mannitol, methyl cellulose, lactose, surfactants such as Tweens or Myrj, magnesium stearate, aqueous or non-aqueous vehicles, paraffin derivatives, cleaning agents, dispersing agents, emulsifiers, preservatives and flavoring substances for taste correction (e.g., ethereal oils).

35

40

The invention thus also relates to pharmaceutical compositions that as active ingredients contain at least one compound according to the invention. A dosage unit will preferably contain about 0.1-100 mg of active ingredient(s). In humans, the dosage of the compounds according to the invention is preferably approximately 0.1-1000 mg per day.

45

50

The examples below are used for a more detailed explanation of the invention, without intending that it be limited to these examples.

55

5

10

15

20

25

30

35

40

45

50

55

The entire disclosure of all applications, patents and publications, cited above or below, and of corresponding German application Nos. 199 21 086.1, filed 30 April 1999, and 199 54 228.7, filed 4 November 1999, are hereby incorporated by reference.

5

107

10

Examples

15

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

20

25

In the foregoing and in the following examples, all temperatures are set forth uncorrected in degrees Celsius; and, unless otherwise indicated, all parts and percentages are by weight.

30

Example 1

35

4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

40

Production of (4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-hydroxy-5-(tetrahydro-2H-pyran-2-yloxy)-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane; Variant I:

45

50

55

5

108

10

Example 1a

(3RS,4S)-4-(2-Methyl-3-hydroxy-8-(trimethylsilyl)-oct-7-in-2-yl)-
2,2-dimethyl-[1,3]dioxane

15

The solution of 6.33 g (34 mmol) of (4S)-4-(2-methyl-1-oxo-prop-2-yl)-2,2-dimethyl-[1,3]dioxane, which was produced analogously to the process described in DE 197 51 200.3, in 10 ml of anhydrous tetrahydrofuran is mixed in portions under an atmosphere of dry argon with the solution of a total of 50 mmol of 5-trimethylsilyl-pent-4-in-1-yl-magnesium bromide in tetrahydrofuran, allowed to heat to 60°C and stirred for 1.5 hours. It is poured onto water and extracted several times with ethyl acetate. The combined organic extracts are washed with water and saturated sodium chloride solution and dried on sodium sulfate. The residue that was obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 6.22 g (19 mmol, 56%) of the chromatographically separable 3R- and 3S-epimers of the title compound and 1.35 g of (4S)-4-(2-methyl-1-hydroxy-prop-2-yl)-2,2-dimethyl-[1,3]dioxane are isolated in each case as a colorless oil.

35

40

45

¹H-NMR (CDCl₃): δ = 0.14 (9H), 0.73+0.88 (3H), 0.91 (3H), 1.28-1.93 (12H), 2.21-2.33 (2H), 3.40-3.72 (2H), 3.80-4.03 (3H) ppm.

50

55

5

109

Example 1b

10

(4S)-4-(2-Methyl-3-oxo-8-(trimethylsilyl)-oct-7-in-2-yl)-2,2-dimethyl-[1,3]dioxane

15

The solution of 6.22 g (19 mmol) of a mixture of the compounds, produced according to Example 1a, in 200 ml of anhydrous dichloromethane is mixed with a molecular sieve (4A, about 20 pellets), 4.01 g of N-methylmorpholino-N-oxide, 335 mg of tetrapropylammonium perruthenate and stirred for 16 hours at 23°C under an atmosphere of dry argon. It is concentrated by evaporation, the crude product that is obtained is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 5.17 g (15.9 mmol, 84%) of the title compound is isolated as a colorless oil.

20

25

30

35

¹H-NMR (CDCl₃): δ = 0.15 (9H), 1.07 (3H), 1.13 (3H), 1.28-1.36 (1H), 1.33 (3H), 1.41 (3H), 1.53-1.81 (3H), 2.22 (2H), 2.62 (2H), 3.85 (1H), 3.97 (1H), 4.06 (1H) ppm.

40

Example 1c

45

(4S(4R,5S,6S,10RS))-4-(5-Hydroxy-2,6-dimethyl-3-oxo-4-(4-(trimethylsilyl)-but-3-in-1-yl)-10-[[diphenyl(1,1-dimethylethyl)silyl]oxy]-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and (4S(4S,5R,6S,10RS))-4-(5-hydroxy-2,6-dimethyl-3-oxo-4-(4-(trimethylsilyl)-but-3-in-1-yl)-10-

50

55

5
10
[[diphenyl(1,1-dimethylethyl)silyl]oxy]-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

15 The solution of 1.33 ml of diisopropylamine in 35 ml of anhydrous tetrahydrofuran is cooled under an atmosphere of dry argon to -30°C, mixed with 4.28 ml of a 2.4 molar solution of n-butyllithium in n-hexane and stirred for another 15 minutes. At
20 -78°C, the solution of 2.87 g (8.84 mmol) of the compound, produced according to Example 1c, in 35 ml of tetrahydrofuran is added in drops, and it is allowed to react for 1 hour. Then, it
25 is mixed slowly with the solution of 3.93 g (10.3 mmol) of (2S,6RS)-2-methyl-6-(tert-butyl-diphenylsilyloxy)-heptanal, which was produced analogously to the process that is described in DE
30 197 51 200.3, in 35 ml of tetrahydrofuran, and it is poured into saturated ammonium chloride solution after 1 hour. It is diluted with water, extracted several times with ethyl acetate, the
35 combined organic extracts are washed with saturated sodium chloride solution, dried on sodium sulfate and concentrated by
40 evaporation in a vacuum. After column chromatography on silica gel with a gradient system that consists of n-hexane and ethyl acetate, 2.40 g (3.39 mmol, 38%) of title compound A and 1.52 g
45 (2.15 mmol, 24%) of diastereomer B are obtained in addition to starting material.

5

111

10

15

¹H-NMR (CDCl₃) of A: δ = 0.16 (9H), 0.83 (3H), 1.00 (3H), 1.02 (3H), 1.04 (9H), 1.10-1.77 (10H), 1.28 (3H), 1.31 (3H), 1.37 (3H), 1.83-2.03 (2H), 2.19-2.38 (2H), 3.52 (1H), 3.62 (1H), 3.78-3.92 (2H), 3.98 (1H), 4.23 (1H), 7.30-7.46 (6H), 7.67 (4H) ppm.

20

¹H-NMR (CDCl₃) of B: δ = 0.13 (9H), 0.86+0.92 (3H), 0.95-1.77 (16H), 1.03 (9H), 1.21+1.25 (3H), 1.32 (3H), 1.40 (3H), 1.88-2.09 (2H), 2.26 (1H), 2.39 (1H), 3.29-3.54 (2H), 3.77-3.90 (2H), 3.96 (1H), 4.18 (1H), 7.31-7.46 (6H), 7.67 (4H) ppm.

25

Example 1d

30

(4S(4R,5S,6S,10RS))-4-(2,6-Dimethyl-3-oxo-4-(4-(trimethylsilyl)-5-(tetrahydro-2H-pyran-2-yloxy)-but-3-in-1-yl)-10-[[diphenyl(1,1-dimethylethyl)silyl]oxy]-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

35

40

45

50

The solution of 2.35 g (3.32 mmol) of compound A, produced according to Example 1c, in 55 ml of anhydrous dichloromethane is mixed under an atmosphere of dry argon with 3.04 ml of 3,4-dihydro-2H-pyran, 0.67 g of p-toluenesulfonic acid, and it is stirred for 48 hours at 23°C. It is poured into a saturated sodium bicarbonate solution, the organic phase is separated, and it is dried on sodium sulfate. After filtration and removal of the solvent, the residue is chromatographed on fine silica gel with a mixture that consists of n-hexane and ethyl acetate. 2.29

55

g (2.89 mmol, 87%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.05 (9H), 0.88-2.15 (28H), 1.03 (9H), 1.41 (3H), 1.59 (3H), 2.21-2.48 (1H), 3.31-4.53 (9H), 7.30-7.45 (6H), 7.69 (4H) ppm.

Example 1e

(4S(4R,5S,6S,10RS))-4-(2,6-Dimethyl-10-hydroxy-3-oxo-5-(tetrahydro-2H-pyran-2-yloxy)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

The solution of 2.48 g (3.13 mmol) of the compound, produced according to Example 1d, in 25 ml of anhydrous tetrahydrofuran is mixed under an atmosphere of dry argon with 12.5 ml of a 1 molar solution of tetrabutylammonium fluoride in tetrahydrofuran, and it is stirred for 4 hours at 23°C. It is mixed with saturated sodium bicarbonate solution, extracted several times with ethyl acetate, washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 1.41 g (2.93 mmol, 94%) of the title compound is isolated as a colorless oil.

Example 1f

(4S(4R,5S,6S,10RS))-4-(2,6-Dimethyl-3,10-dioxo-5-(tetrahydro-2H-pyran-2-yloxy)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

Analogously to Example 1b, 1.27 g (2.63 mmol) of the compound, produced according to Example 1e, is reacted, and after working-up and purification, 1.14 g (2.38 mmol, 91%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.95-2.48 (29H), 0.98+1.01 (3H), 1.42 (3H), 2.13 (3H), 3.29-3.47 (2H), 3.64-4.04 (4H), 4.20+4.32 (1H), 4.39+4.50 (1H) ppm.

Example 1g

(4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-[[Diphenyl(1,1-dimethylethyl)silyl]oxy]-5-(tetrahydro-2H-pyran-2-yloxy)-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

The suspension of 2.87 g (3.57 mmol) of (5E,3S)-[3-[[[(1,1-dimethylethyl)diphenylsilyl]oxy]-4-methyl-5-(2-pyridyl)-pent-4-en-1-yl]-triphenyl-phosphonium iodide, which was produced analogously to the process described in DE 197 51 200.3, in 11 ml of anhydrous tetrahydrofuran, is mixed at 0°C under an atmosphere of dry argon with 2.72 ml of a 1.6 M solution of n-butyllithium

in n-hexane and allowed to heat to 23°C. The solution of 1.14 g (2.38 mmol) of the compound, produced according to Example 1f, in 11 ml of tetrahydrofuran is slowly added in drops to the red solution, allowed to stir for 2 hours, poured onto saturated ammonium chloride solution and extracted several times with ethyl acetate. The combined organic extracts are dried on sodium sulfate and concentrated by evaporation in a vacuum. After column chromatography on silica gel with a gradient system that consists of n-hexane and ethyl acetate, 860 mg (0.98 mmol, 41%) of the title compound is obtained in addition to 20% starting material.

¹H-NMR (CDCl₃): δ = 0.82-2.41 (41H), 1.05 (9H), 2.00 (3H), 3.23-3.45 (2H), 3.60-4.02 (3H), 4.08-4.51 (3H), 4.92-5.24 (1H), 6.16-6.76 (1H), 6.92-7.08 (2H), 7.21-7.43 (6H), 7.49-7.72 (5H), 8.55 (1H) ppm.

Example 1h; Variant I: (4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-Hydroxy-5-(tetrahydro-2H-pyran-2-yloxy)-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

Analogously to Example 1b, 482 mg (550 μmol) of the compound that is produced according to Example 1g is reacted, and after

working-up and purification, 256 mg (401 μ mol, 73%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.88-2.48 (35H), 1.42 (3H), 1.64+1.72 (3H), 2.08 (3H), 3.29-3.47 (2H), 3.64-4.04 (4H), 4.12-4.35 (2H), 4.41+4.51 (1H), 5.20 (1H), 6.59 (1H), 7.09 (1H), 7.23 (1H), 7.63 (1H), 8.60 (1H) ppm.

Production of (4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-hydroxy-5-(tetrahydro-2H-pyran-2-yloxy)-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane; Variant II:

Example 1i

(4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-[[Diphenyl(1,1-dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(4-(trimethylsilyl)-but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and
(4S(4S,5R,6S,10E/Z,13S,14E))-4-(13-[[diphenyl(1,1-dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(4-(trimethylsilyl)-but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

Analogously to Example 1c, 2.85 g (8.78 mmol) of the compound, produced according to Example 1b, is reacted with 3.62

g (6.71 mmol) of (2S,6E/Z,9S,10E)-2,6,10-trimethyl-9-
[[diphenyl(1,1-dimethylethyl)silyl]oxy]-1-oxo-11-(2-pyridyl)-
undeca-6,10-diene, which was produced analogously to the process
that is described in DE 197 51 200.3, and after working-up and
purification, in addition to starting material, 1.28 g (1.48
mmol, 22%) of title compound C as well as 1.73 g (2.00 mmol, 30%)
of title compound B are isolated in each case as a colorless oil.

¹H-NMR (CDCl₃) of A: δ = 0.13 (9H), 0.86-2.52 (36H), 1.08
(9H), 1.42+1.58 (3H), 2.01 (3H), 3.32-4.85 (9H), 5.00 (1H), 6.23
(1H), 6.97-7.09 (2H), 7.21-7.45 (6H), 7.57 (1H), 7.61-7.75 (4H),
8.56 (1H) ppm.

¹H-NMR (CDCl₃) of B: δ = 0.12 (9H), 0.77-2.53 (36H), 1.08
(9H), 1.38+1.62 (3H), 2.00 (3H), 3.23-4.86 (9H), 5.02 (1H), 6.23
(1H), 6.96-7.09 (2H), 7.19-7.47 (6H), 7.53-7.76 (5H), 8.57 (1H)
ppm.

Example 1j

(4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-[[Diphenyl(1,1-
dimethylethyl)silyl]oxy]-5-(tetrahydro-2H-pyran-2-yloxy)-
2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(4-(trimethylsilyl)-
but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

Analogously to Example 1d, 1.16 g (1.34 mmol) of the
compound that is produced according to Example 1i is reacted, and

5

117

10

after working-up and purification, 1.12 g (1.18 mmol, 88%) of the title compound is isolated as a colorless oil.

15

¹H-NMR (CDCl₃): δ = 0.13 (9H), 0.86-2.52 (39H), 1.08 (9H), 2.01 (3H), 3.32-4.85 (9H), 5.00 (1H), 6.22 (1H), 6.96-7.09 (2H), 7.21-7.44 (6H), 7.56 (1H), 7.61-7.75 (4H), 8.56 (1H) ppm.

20

Example 1h; Variant II (4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-Hydroxy-5-(tetrahydro-2H)-pyran-2-yloxy)-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(but-3-in-1-yl)-undec-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

25

30

Analogously to Example 1e, 1.12 g (1.18 mmol) of the compound that is produced according to Example 1j is reacted, and after working-up and purification, 654 mg (1.03 mmol, 87%) of the title compound is isolated as a colorless oil. The coverage of the ¹H-NMR spectrum is identical to that described in Example 1h, Variant I.

35

40

Example 1k

45

(3S,6R,7S,8S,12E/Z,15S,16E)-1,3,7,15-Tetrahydroxy-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dien-5-one

50

The solution of 654 mg (1.03 mmol) of the compound, produced according to Example 1h, in 27 ml of anhydrous ethanol is mixed

55

5

118

10

15

under an atmosphere of dry argon with 588 mg of p-toluenesulfonic acid-monohydrate, and it is stirred for 3 hours at 23°C. After removal of the solvent, the residue is chromatographed on fine silica gel with a mixture of n-hexane and ethyl acetate. 484 mg (942 μ mol, 91%) of the title compound is isolated as a colorless oil.

20

25

$^1\text{H-NMR}$ (CDCl_3): δ = 0.90+0.92 (3H), 1.07 (3H), 1.11-2.16 (14H), 1.29 (3H), 1.63+1.42 (3H), 2.00+2.02 (3H), 2.20-2.60 (4H), 2.98 (1H), 3.48-3.67 (2H), 3.78-3.93 (2H), 4.06-4.23 (3H), 5.16+5.24 (1H), 6.52+6.57 (1H), 7.11 (1H), 7.30 (1H), 7.66 (1H), 8.58 (1H) ppm.

30

Example 11

35

(3S,6R,7S,8S,12E/Z,15S,16E)-1,3,7,15-Tetrakis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dien-5-one

40

45

50

The solution of 673 mg (1.31 mmol) of the compound, produced according to Example 1k, in 37 ml of anhydrous dichloromethane is cooled under an atmosphere of dry argon to -78°C, mixed with 2.14 ml of 2,6-lutidine, 2.41 ml of trifluoromethanesulfonic acid-tert-butyldimethylsilylester, allowed to heat within 2 hours to 0°C and stirred for another 2 hours. It is poured into saturated sodium bicarbonate solution and extracted several times with

55

5

119

10

dichloromethane. The combined organic extracts are dried on sodium sulfate and concentrated by evaporation in a vacuum. After column chromatography on silica gel with a gradient system that consists of n-hexane and ethyl acetate, 1.11 g (1.29 mmol, 99%) of the title compound is isolated as a colorless oil.

15

20

$^1\text{H-NMR}$ (CDCl_3): δ = -0.01-0.12 (24H), 0.82-2.33 (55H), 1.08 (3H), 1.22 (3H), 1.60+1.68 (3H), 2.05 (3H), 3.22 (1H), 3.51-3.73 (2H), 3.81 (1H), 3.92 (1H), 4.11 (1H), 5.18 (1H), 6.47 (1H), 7.08 (1H), 7.22 (1H), 7.61 (1H), 8.59 (1H) ppm.

25

Example 1m

30

(3S,6R,7S,8S,12E/Z,15S,16E)-1-Hydroxy-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dien-5-one

35

40

The solution of 1.10 mg (1.13 mmol) of the compound, produced according to Example 1l, in a mixture of 14 ml of dichloromethane and 14 ml of methanol is mixed at 23°C under an atmosphere of dry argon with 312 mg of campher-10-sulfonic acid, and it is stirred for 2 hours. It is poured into a saturated sodium bicarbonate solution and extracted several times with dichloromethane. The combined organic extracts are dried on sodium sulfate and concentrated by evaporation in a vacuum.

45

50

After column chromatography on fine silica gel with a gradient

55

system that consists of n-hexane and ethyl acetate, 814 mg (950 μ mol, 84%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.01-0.13 (18H), 0.83-2.33 (47H), 1.12 (3H), 1.23 (3H), 1.61+1.68 (3H), 2.05 (3H), 3.28 (1H), 3.68 (2H), 3.84 (1H), 4.02-4.18 (2H), 5.18 (1H), 6.48 (1H), 7.08 (1H), 7.22 (1H), 7.61 (1H), 8.60 (1H) ppm.

Example 1n

(3S,6R,7S,8S,12E/Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dienal

The solution of 0.129 ml of oxalyl chloride in 6.3 ml of anhydrous dichloromethane is cooled under an atmosphere of dry argon to -70°C , mixed with 209 μ l of dimethyl sulfoxide, the solution of 814 mg (950 μ mol) of the compound, produced according to Example 1m, in 6.3 ml of anhydrous dichloromethane, and it is stirred for 0.5 hour. Then, it is mixed with 646 μ l of triethylamine, allowed to react for 1 hour at -30°C and mixed with n-hexane and saturated sodium bicarbonate solution. The organic phase is separated, the aqueous phase is extracted several more times with n-hexane, the combined organic extracts are washed with water and dried on magnesium sulfate. The

5

121

10

residue that is obtained after filtration and removal of the solvent is further reacted without purification.

15

Example 10

20

(3S,6R,7S,8S,12Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dienoic acid (A) and (3S,6R,7S,8S,12E,15S,16E)-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dienoic acid (B)

25

30

The solution of 852 mg (max. 950 μ mol) of the compound, produced according to Example 1n, in 23 ml of acetone is cooled to -30°C, mixed with 1.19 ml of a standardized, 8N chromosulfuric acid solution and stirred for 1 hour. It is poured into a mixture that consists of water and diethyl ether, the organic phase is washed with saturated sodium chloride solution and dried on sodium sulfate. After filtration and removal of the solvent, the residue is purified by chromatography on a fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 298 mg (342 μ mol, 36% relative to the educt in Example 11) of title compound A as well as 234 mg (269 μ mol, 28% relative to the educt in Example 11) of title compound B are isolated in each case as a colorless oil.

35

40

45

50

55

5

122

10

¹H-NMR (CDCl₃) of A: δ = -0.02-0.15 (18H), 0.81-0.99 (30H), 1.05-2.3 (15H), 1.12 (3H), 1.24 (3H), 1.71 (3H), 1.92 (3H), 2.38 (1H), 2.51 (1H), 3.27 (1H), 3.80 (1H), 4.17 (1H), 4.43 (1H), 5.23 (1H), 6.67 (1H), 7.18 (1H), 7.36 (1H), 7.72 (1H), 8.62 (1H) ppm.

15

20

¹H-NMR (CDCl₃) of B: δ = -0.01-0.19 (18H), 0.80-0.96 (30H), 1.00-2.45 (16H), 1.13 (3H), 1.27 (3H), 1.57 (3H), 1.94 (3H), 2.54 (1H), 3.28 (1H), 3.88 (1H), 4.13 (1H), 4.40 (1H), 5.12 (1H), 6.49 (1H), 7.18 (1H), 7.38 (1H), 7.71 (1H), 8.62 (1H) ppm.

25

Example 1p

30

(3S,6R,7S,8S,12Z,15S,16E)-15-Hydroxy-3,7-bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dienoic acid

35

Analogously to Example 1e, 298 mg (342 μmol) of compound A, produced according to Example 1o, is reacted, and after working-up, 294 mg (max. 342 μmol) of the title compound is isolated as a crude product, which is further reacted without purification.

40

Example 1q

45

(4S,7R,8S,9S,13Z,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

50

55

5

123

10

15

20

25

30

35

40

The solution of 294 mg (max. 342 μ mol) of the compound, produced according to Example 1p, in a mixture that consists of 2.6 ml of anhydrous tetrahydrofuran and 30 ml of toluene is mixed under an atmosphere of dry argon with 284 μ l of triethylamine, 268 μ l of 2,4,6-trichlorobenzoyl chloride, and it is stirred for 20 minutes. This solution is added in drops within 4.5 hours to the solution of 434 mg of 4-dimethylaminopyridine in 132 ml of toluene, and it is stirred for 0.5 more hour at 23°C. It is concentrated by evaporation, taken up in a little dichloromethane and purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 136 mg (184 μ mol, 54%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = -0.08 (3H), 0.13 (9H), 0.80-2.32 (12H), 0.85 (9H), 0.94 (9H), 0.99 (3H), 1.15 (3H), 1.24 (3H), 1.68 (3H), 2.13 (3H), 2.47 (1H), 2.59-2.89 (3H), 3.11 (1H), 4.00 (1H), 4.06 (1H), 5.00 (1H), 5.18 (1H), 6.57 (1H), 7.10 (1H), 7.26 (1H), 7.63 (1H), 8.60 (1H) ppm.

Example 1r

45

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

50

55

The solution of 20 mg (27 μ mol) of the compound, produced according to Example 1p, in 2 ml of anhydrous tetrahydrofuran is mixed under an atmosphere of dry argon in portions with a total of 0.57 ml of HF-pyridine complex, and it is stirred at 23°C for 24 hours. It is poured into saturated sodium bicarbonate solution, extracted several times with dichloromethane, and the combined organic extracts are dried on sodium sulfate. After filtration and removal of the solvent, the residue that is obtained is purified by chromatography on fine silica gel with a mixture that consists of n-hexane and ethyl acetate. 9.1 mg (17.9 μ mol, 66%) of the title compound is isolated as a colorless oil as well as mg of monosilylether.

¹H-NMR (CDCl₃): δ = 1.09 (6H), 1.19-2.12 (11H), 1.38 (3H), 6.9 (3H), 2.06 (3H), 2.21-2.41 (3H), 2.50 (1H), 2.63 (1H), 2.68 (1H), 3.53 (1H), 3.70 (1H), 4.42 (1H), 4.59 (1H), 5.12 (1H), 5.22 (1H), 6.61 (1H), 7.13 (1H), 7.29 (1H), 7.68 (1H), 8.53 (1H) ppm.

Example 2

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Example 2a

(4S,7R,8S,9S,13Z,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

The solution of 25 mg (34 μ mol) of the compound, produced according to Example 1q, in 3 ml of ethanol, is mixed with 25 μ l of pyridine, a catalytic amount of palladium on barium sulfate (10%), and it is hydrogenated under 1 atmosphere of hydrogen. After filtration and removal of the solvent, the residue is purified by chromatography on an analytical thin-layer plate. As a mobile solvent, a mixture of n-hexane and ethyl acetate is used. 13 mg (18 μ mol, 52%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = -0.10 (3H), 0.06 (3H), 0.11 (6H), 0.80-2.20 (11H), 0.83 (9H), 0.92 (9H), 0.98 (3H), 1.12 (3H), 1.19 (3H), 1.67 (3H), 2.12 (3H), 2.43 (1H), 2.55-2.82 (3H), 3.07 (1H), 4.00 (1H), 4.03 (1H), 4.90-5.03 (3H), 5.18 (1H), 5.72 (1H), 6.57 (1H), 7.09 (1H), 7.25 (1H), 7.62 (1H), 8.59 (1H) ppm.

Example 2b

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 1, 10.3 mg (14 μ mol) of the compound that is produced according to Example 2a is reacted, and after working-up and purification, 5.7 mg (11 μ mol, 80%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 1.04 (3H), 1.09 (3H), 0.25-2.38 (13H), 1.36 (3H), 1.70 (3H), 2.07 (3H), 2.48 (1H), 2.63 (1H), 2.74 (1H), 3.31 (1H), 3.69 (1H), 4.38 (1H), 4.61 (1H), 4.97 (1H), 5.02 (1H), 5.11 (1H), 5.19 (1H), 5.77 (1H), 6.60 (1H), 7.13 (1H), 7.29 (1H), 7.68 (1H), 8.54 (1H) ppm.

Example 3

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

Example 3a

(3S,6R,7S,8S,12E,15S,16E)-15-Hydroxy-3,7-bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(but-3-in-1-yl)-heptadeca-12,16-dienoic acid

Analogously to Example 1e, 234 mg (269 μ mol) of compound B that is produced according to Example 1o is reacted, and after working-up, 229 mg (max. 269 μ mol) of the title compound is isolated as a crude product, which is further reacted without purification.

Example 3b

(4S,7R,8S,9S,13E,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 1q, 229 mg (max. 269 μ mol) of the compound that is produced according to Example 3a is reacted, and after working-up and purification, 112 mg (152 μ mol, 56%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.05 (3H), 0.11 (6H), 0.15 (3H), 0.80-2.30 (33H), 1.13 (3H), 1.21 (3H), 1.62 (3H), 2.61 (3H), 2.40-2.72 (4H), 3.10 (1H), 3.91 (1H), 4.46 (1H), 5.22 (1H), 5.30 (1H), 6.56 (1H), 7.09 (1H), 7.20 (1H), 7.62 (1H), 8.60 (1H) ppm.

5

128

10

Example 3c

15

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

20

Analogously to Example 1, 72 mg (98 μ mol) of the compound that is produced according to Example 3b is reacted, and after working-up and purification, 32 mg (63 μ mol, 64%) of the title compound is isolated as a colorless foam.

25

$^1\text{H-NMR}$ (CDCl_3): δ = 1.00 (3H), 1.04 (3H), 1.30-2.71 (16H), 1.32 (3H), 1.61 (3H), 2.10 (3H), 3.63 (1H), 3.70 (1H), 3.86 (1H), 3.99 (1H), 4.48 (1H), 5.10 (1H), 5.41 (1H), 6.58 (1H), 7.13 (1H), 7.33 (1H), 7.68 (1H), 8.54 (1H) ppm.

30

Example 4

35

(1S,3S(E),7S,10R,11S,12S,16R)-7,11-Dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and

40

(1R,3S(E),7S,10R,11S,12S,16S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

45

50

The solution of 5 mg (10 μ mol) of the compound, produced according to Example 1, in 1 ml of dichloromethane is mixed under an atmosphere of dry argon at -20°C with 11.3 μ l of a 20%

55

5

129

10

solution of trifluoroacetic acid in dichloromethane and 5.6 mg of m-chloroperbenzoic acid (60%). It is stirred for 18 hours at -18°C, poured onto saturated sodium thiosulfate solution,

15

extracted several times with dichloromethane, the combined organic extracts are washed with sodium bicarbonate solution, saturated sodium chloride solution and dried on magnesium

20

sulfate. The residue that is obtained after filtration and

removal of the solvent is purified by chromatography on an analytical thin-layer plate. As a mobile solvent and eluant, a mixture that consists of dichloromethane and ethanol is used.

25

1.3 mg (2.5 μ mol, 25%) of title compound A (or B) and 2.0 mg (3.8 μ mol, 39%) of title compound B (or A) are isolated in each case as a colorless oil.

30

¹H-NMR (CDCl₃) of A (or B): δ = 1.01 (3H), 1.07 (3H), 1.23-

35

2.20 (13H), 1.30 (3H), 1.46 (3H), 2.10 (3H), 2.26 (1H), 2.40 (1H), 2.58 (1H), 2.82 (1H), 2.97 (1H), 3.63 (2H), 4.39 (1H), 5.22 (1H), 5.47 (1H), 6.61 (1H), 7.15 (1H), 7.28 (1H), 7.69 (1H), 8.55 (1H) ppm.

40

¹H-NMR (CDCl₃) of B (or A): δ = 0.98 (3H), 1.08 (3H), 1.27-

45

2.19 (13H), 1.32 (3H), 1.43 (3H), 2.12 (3H), 2.30 (1H), 2.48 (1H), 2.70 (1H), 2.96 (1H), 3.15 (1H), 3.47 (1H), 3.57 (1H), 4.01 (1H), 4.49 (1H), 5.50 (1H), 6.67 (1H), 7.12 (1H), 7.27 (1H), 7.66 (1H), 8.58 (1H) ppm.

50

55

Example 5

(1S,3S(E),7S,10R,11S,12S,16R)-7,11-Dihydroxy-10-(but-3-en-1-yl)-
3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(E),7S,10R,11S,12S,16S)-7,11-dihydroxy-10-(but-3-en-1-yl)-
3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 4, 6.6 mg (13 μ mol) of the compound that is produced according to Example 2 is reacted, and after working-up and purification, 1.4 mg (2.7 μ mol, 20%) of title compound A (or B) and 0.9 mg (1.7 μ mol, 13%) of title compound B (or A) are isolated in each case as a colorless foam.

$^1\text{H-NMR}$ (CDCl_3) of A (or B): δ = 1.00 (3H), 1.07 (3H), 1.21-2.05 (12H), 1.30 (3H), 1.40 (3H), 2.10 (3H), 2.16 (1H), 2.38 (1H), 2.57 (1H), 2.81 (1H), 2.97 (1H), 3.44 (1H), 3.63 (1H), 4.38 (1H), 4.98 (1H), 5.02 (1H), 5.28 (1H), 5.45 (1H), 5.77 (1H), 6.62 (1H), 7.18 (1H), 7.31 (1H), 7.71 (1H), 8.56 (1H) ppm.

$^1\text{H-NMR}$ (CDCl_3) of B (or A): δ = 0.94 (3H), 1.05 (3H), 1.18-2.17 (13H), 1.30 (3H), 1.38 (3H), 2.12 (3H), 2.48 (1H), 2.62 (1H), 2.95 (1H), 3.28 (1H), 3.30 (1H), 3.50 (1H), 3.96 (1H), 4.41 (1H), 4.95 (1H), 5.00 (1H), 5.52 (1H), 5.25 (1H), 6.73 (1H), 7.18 (1H), 7.33 (1H), 7.71 (1H), 8.58 (1H) ppm.

Example 6

(1S,3S(E),7S,10R,11S,12S,16S)-7,11-Dihydroxy-10-(but-3-in-1-yl)-
3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(E),7S,10R,11S,12S,16R)-7,11-dihydroxy-10-(but-3-in-1-yl)-
3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 4, 14 mg (27 μ mol) of the compound that is produced according to Example 3 is reacted, and after working-up and purification, 7.8 mg (15 mol, 55%) of title compound A (or B) and 4.7 mg (9 μ mol, 33%) of title compound B (or A) are isolated in each case as a colorless foam.

$^1\text{H-NMR}$ (CDCl_3) of A (or B): δ = 0.93 (3H), 1.04 (3H), 1.23-2.19 (13H), 1.29 (3H), 1.42 (3H), 2.13 (3H), 2.28 (1H), 2.48-2.65 (2H), 2.71 (1H), 2.89 (1H), 3.57 (1H), 3.83 (1H), 4.36 (1H), 4.47 (1H), 5.51 (1H), 6.63 (1H), 7.12 (1H), 7.28 (1H), 7.67 (1H), 8.57 (1H) ppm.

$^1\text{H-NMR}$ (CDCl_3) of B (or A): δ = 0.96 (3H), 1.10 (3H), 1.21-2.18 (13H), 1.26 (3H), 1.40 (3H), 2.10 (3H), 2.29 (1H), 2.61 (2H), 2.86 (1H), 2.99 (1H), 3.58 (1H), 3.79 (2H), 4.37 (1H), 5.46 (1H), 6.61 (1H), 7.12 (1H), 7.26 (1H), 7.66 (1H), 8.57 (1H) ppm.

Example 7

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 2a, 14 mg (27 μ mol) of the compound that is produced according to Example 3 is reacted, and after working-up and purification, 4.1 mg (8 μ mol, 29%) of the title compound is isolated as a colorless foam.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.98 (3H), 1.02 (3H), 1.30 (3H), 1.36-2.68 (16H), 1.61 (3H), 2.09 (3H), 3.43 (1H), 3.70 (1H), 4.17 (1H), 4.45 (1H), 4.94 (1H), 5.00 (1H), 5.09 (1H), 5.39 (1H), 5.72 (1H), 6.58 (1H), 7.12 (1H), 7.35 (1H), 7.67 (1H), 8.52 (1H) ppm.

Example 8

(1S,3S(E),7S,10R,11S,12S,16S)-7,11-Dihydroxy-10-(but-3-en-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(E),7S,10R,11S,12S,16R)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 4, 4.1 mg (8 μ mol) of the compound that is produced according to Example 7 is reacted, and after working-up and purification, 1.7 mg (3.2 μ mol, 40%) of title

5

10

compound A (or B) and 0.4 mg (0.8 μ mol, 9%) of title compound B (or A) are isolated in each case as a colorless foam.

15

$^1\text{H-NMR}$ (CDCl_3) of A (or B): δ = 0.91 (3H), 1.02 (3H), 1.13-2.17 (15H), 1.28 (3H), 1.38 (3H), 2.11 (3H), 2.53 (2H), 2.87 (1H), 2.96 (1H), 3.38 (1H), 3.78 (1H), 4.35 (1H), 4.37 (1H), 4.95 (1H), 5.00 (1H), 5.50 (1H), 5.76 (1H), 6.64 (1H), 7.12 (1H), 7.30 (1H), 7.67 (1H), 8.57 (1H) ppm.

20

25

$^1\text{H-NMR}$ (CDCl_3) of B (or A): δ = 0.92 (3H), 1.09 (3H), 1.18-2.13 (15H), 1.26 (3H), 1.38 (3H), 2.08 (3H), 2.49-2.60 (2H), 2.85-2.99 (2H), 3.39 (1H), 3.72 (1H), 3.89 (1H), 4.28 (1H), 4.92-5.06 (2H), 5.45 (1H), 5.76 (1H), 6.60 (1H), 7.12 (1H), 7.26 (1H), 7.68 (1H), 8.57 (1H) ppm.

30

Example 9

35

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

40

Example 9a

45

(3RS,4S)-4-(2-Methyl-3-hydroxy-hept-6-en-2-yl)-2,2-dimethyl-[1,3]dioxane

50

Analogously to Example 1a, 5.5 g (30 mmol) of (4S)-4-(2-methyl-1-oxo-prop-2-yl)-2,2-dimethyl-[1,3]dioxane, which was

55

5

10

15

produced analogously to the process described in DE 197 51 200.3,
is reacted with but-3-en-1-yl-magnesium bromide, and after
working-up and purification, 3.84 g (15.8 mmol, 53%) of the title
compound is isolated as a colorless oil.

Example 9b

20

(4S)-4-(2-Methyl-3-oxo-hept-6-en-2-yl)-2,2-dimethyl-[1,3]dioxane

25

Analogously to Example 1b, 3.84 g (15.8 mmol) of the
compound that is produced according to Example 9a is reacted, and
after working-up and purification, 3.0 g (12.5 mmol, 79%) of the
title compound is isolated as a colorless oil.

30

¹H-NMR (CDCl₃): δ = 1.07 (3H), 1.14 (3H), 1.33 (4H), 1.41
(3H), 1.62 (1H), 2.29 (2H), 2.60 (2H), 3.86 (1H), 3.97 (1H), 4.05
(1H), 4.96 (1H), 5.02 (1H), 5.81 (1H) ppm.

35

Example 9c

40

(4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-[[Dimethyl(1,1-

45

dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-tetramethyl-3-oxo-
15-(2-methylthiazol-4-yl)-4-(prop-2-en-1-yl)-pentadec-2-yl)-2,2-
dimethyl-[1,3]dioxane (A) and (4S(4S,5R,6S,10E/Z,13S,14E))-4-(13-
[[dimethyl(1,1-dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-
tetramethyl-3-oxo-15-(2-methylthiazol-4-yl)-4-(prop-2-en-1-yl)-
pentadec-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

50

55

Analogously to Example 1c, 2.07 g (8.61 mmol) of the compound, produced according to Example 9b, with 2.01 g (4.61 mmol) of (2S,6E/Z,9S,10E)-2,6,10-trimethyl-9-[dimethyl(1,1-dimethylethyl)silyl]oxy]-1-oxo-11-(2-methylthiazol-4-yl)-undeca-6,10-diene that was produced analogously to the process that is described in DE 197 51 200.3 is reacted, and after working-up and purification, in addition to starting material, 995 mg (1.47 mmol, 32%) of title compound A as well as 784 mg (1.16 mmol, 25%) of title compound B are isolated in each case as a colorless oil.

¹H-NMR (CDCl₃) of A: δ = 0.01 (3H), 0.07 (3H), 0.85 (3H), 0.90 (9H), 0.98 (3H), 1.00-2.33 (12H), 1.23 (3H), 1.33 (3H), 1.39 (3H), 1.60+1.67 (3H), 2.00 (3H), 2.46 (1H), 2.72 (3H), 2.99 (1H), 3.34 (1H), 3.49 (1H), 3.87 (1H), 3.98 (1H), 4.09 (1H), 4.13 (1H), 4.98 (1H), 5.03 (1H), 5.13 (1H), 5.71 (1H), 6.44 (1H), 6.93 (1H) ppm.

¹H-NMR (CDCl₃) of B: δ = 0.00 (3H), 0.03 (3H), 0.88 (9H), 0.94 (3H), 1.03-1.72 (7H), 1.08 (3H), 1.17 (3H), 1.31 (3H), 1.39 (3H), 1.60+1.68 (3H), 1.89-2.08 (2H), 1.99 (3H), 2.17-2.51 (4H), 2.71 (3H), 2.74+2.87 (1H), 3.31 (1H), 3.57 (1H), 3.84 (1H), 3.95 (1H), 4.03-4.17 (2H), 4.98 (1H), 5.03 (1H), 5.13 (1H), 5.73 (1H), 6.64 (1H), 6.92 (1H) ppm.

5

Example 9d

10

(3S,6R,7S,8S,12E/Z,15S,16E)-15-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-1,3,7-trihydroxy-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-6-(prop-2-en-1-yl)-heptadeca-12,16-dien-5-one

15

Analogously to Example 1k, 1.33 g (1.97 mmol) of compound A that is produced according to Example 9c is reacted, and after working-up and purification, 1.02 g (1.60 mmol, 81%) of the title compound is isolated as a colorless oil.

20

25

¹H-NMR (CDCl₃): δ = 0.01 (3H), 0.07 (3H), 0.89 (12H), 1.00-2.38 (12H), 1.40+1.07 (3H), 1.23+1.25 (3H), 1.60+1.68 (3H), 1.97+1.99 (3H), 2.52 (1H), 2.67-2.89 (1H), 2.73+2.77 (3H), 3.01 (1H), 3.33 (1H), 3.40-3.53 (1H), 3.74-3.93 (3H), 4.03-4.19 (2H), 5.00 (1H), 5.06 (1H), 5.10+5.20 (1H), 5.71 (1H), 6.42 (1H), 6.93 (1H) ppm.

30

35

Example 9e

40

(3S,6R,7S,8S,12E/Z,15S,16E)-1,3,7,15-Tetrakis-[[dimethyl(1,1-dimethylethyl)silyl]oxy[4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-6-(prop-2-en-1-yl)-heptadeca-12,16-dien-5-one

45

Analogously to Example 1l, 1.02 g (1.60 mmol) of the compound that is produced according to Example 9d is reacted, and

50

55

after working-up and purification, 1.46 g (1.49 mmol, 93%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.00-0.11 (24H), 0.83-0.98 (39H), 1.01-1.62 (8H), 1.07 (3H), 1.20 (3H), 1.59+1.67 (3H), 1.97 (1H), 2.00 (3H), 2.19-2.34 (3H), 2.48 (1H), 2.72 (3H); 3.13 (1H), 3.57 (1H), 3.67 (1H), 3.78 (1H), 3.87 (1H), 4.09 (1H), 4.93 (1H), 4.99 (1H), 5.15 (1H), 5.77 (1H), 6.64 (1H), 6.91 (1H) ppm.

Example 9f

(3S,6R,7S,8S,12E/Z,15S,16E)-1-Hydroxy-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-6-(prop-2-en-1-yl)-heptadeca-12,16-dien-5-one

Analogously to Example 1m, 1.45 g (1.48 mmol) of the compound that is produced according to Example 9e is reacted, and after working-up and purification, 1.19 g (1.37 mmol, 93%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.01-0.14 (18H), 0.82-0.97 (30H), 1.04-1.70 (7H), 1.09 (3H), 1.19 (3H), 1.59+1.68 (3H), 1.84-2.08 (3H), 2.00 (3H), 2.18-2.36 (3H), 2.47 (1H), 2.71 (3H), 3.13 (1H), 3.66 (2H), 3.80 (1H), 4.40 (1H), 4.10 (1H), 4.96 (1H), 5.01 (1H), 5.14 (1H), 5.77 (1H), 6.46 (1H), 6.92 (1H) ppm.

5

Example 9g

10

(3S,6R,7S,8S,12E/Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-5-oxo-6-(prop-2-en-1-yl)-heptadeca-12,16-dienal

15

Analogously to Example 1n, 1.18 g (1.37 mmol) of the compound that is produced according to Example 9f is reacted, and after working-up, 1.25 g (max. 1.37 mmol) of the title compound is isolated as a yellow oil, which is further reacted without purification.

25

Example 9h

30

(3S,6R,7S,8S,12Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-5-oxo-6-(prop-2-en-1-yl)-heptadeca-12,16-dienoic acid (A) and (3S,6R,7S,8S,12E,15S,16E)-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-5-oxo-6-(prop-2-en-1-yl)-heptadeca-12,16-dienoic acid (B)

35

40

Analogously to Example 1o, 1.25 g (max. 1.37 mmol) of the compound that is produced according to Example 9g is reacted, and after working-up and purification, 302 mg (0.34 mmol, 25%) of

45

50

55

5

10

title compound A and 230 mg (0.26 mmol, 19%) of title compound B are isolated in each case as a colorless oil.

15

¹H-NMR (CDCl₃) of A: δ = -0.02-0.15 (18H), 0.82-0.97 (30H), 1.05-2.53 (14H), 1.12 (3H), 1.17 (3H), 1.70 (3H), 1.96 (3H), 2.71 (3H), 3.17 (1H), 3.72 (1H), 4.16 (1H), 4.37 (1H), 4.94 (1H), 4.99 (1H), 5.20 (1H), 5.73 (1H), 6.66 (1H), 6.93 (1H) ppm.

20

¹H-NMR (CDCl₃) of B: δ = -0.03-0.15 (18H), 0.81-0.95 (30H), 1.01-2.50 (13H), 1.12 (3H), 1.18 (3H), 1.57 (3H), 1.95 (3H), 2.60 (1H), 2.70 (3H), 3.22 (1H), 3.79 (1H), 4.08 (1H), 4.32 (1H), 4.94 (1H), 5.00 (1H), 5.11 (1H), 5.74 (1H), 6.46 (1H), 6.93 (1H) ppm.

25

Example 9i

30

(3S,6R,7S,8S,12Z,15S,16E)-3,7-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-15-hydroxy-4,4,8,12,16-pentamethyl-17-(2-methylthiazol-4-yl)-5-oxo-6-(prop-2-en-1-yl)-heptadecane-12,16-dienoic acid

35

40

Analogously to Example 1e, 302 mg (0.34 mmol) of compound A that is produced according to Example 9h is reacted, and after working-up, 296 mg (max. 0.34 mmol) of the title compound is isolated as a pale yellow oil, which is further reacted without purification.

45

50

55

5

140

10

Example 9j

15

(4S,7R,8S,9S,13Z,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-methylthiazol-4-yl)-ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

20

Analogously to Example 1q, 296 mg (max. 0.34 mmol) of the compound that is produced according to Example 9i is reacted, and after working-up and purification, 166 mg (0.22 mmol, 65%) of the title compound is isolated as a colorless oil.

25

¹H-NMR (CDCl₃): δ = -0.10 (3H), 0.09 (3H), 0.11 (3H), 0.13 (3H), 0.86 (9H), 0.80-2.85 (13H), 0.94 (9H), 1.00 (3H), 1.10 (3H), 12.0 (3H), 1.68 (3H), 2.10 (3H), 2.71 (3H), 3.11 (1H), 4.01 (2H), 4.85-5.03 (3H), 5.16 (1H), 5.78 (1H), 6.57 (1H), 6.98 (1H) ppm.

35

Example 9k

40

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

Analogously to Example 1, 25 mg (34 μmol) of the compound that is produced according to Example 9j is reacted, and after working-up and purification, 10 mg (19 μmol, 57%) of the title compound is isolated as a colorless oil.

50

55

5

10

15

¹H-NMR (CDCl₃): δ = 1.03 (3H), 1.05 (3H), 1.20-2.74 (14H),
1.30 (3H), 1.69 (3H), 2.07 (3H), 2.69 (3H), 3.33 (1H), 3.69 (1H),
3.72 (1H), 4.23 (1H), 5.02 (1H), 5.07 (1H), 5.12 (1H), 5.21 (1H),
5.76 (1H), 6.57 (1H), 6.96 (1H) ppm.

Example 10

20

25

30

(1S,3S(E),7S,10R,11S,12S,16R)-7,11-Dihydroxy-10-(prop-2-en-1-yl)-
3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A)
and (1R,3S(E),7S,10R,11S,12S,16S)-7,11-dihydroxy-10-(prop-2-en-1-
yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

35

40

45

50

The solution of 8.0 mg (15.5 μmol) of the compound, produced according to Example 9, in 1 ml of acetonitrile is mixed with 89 μl of a 1M solution of sodium ethylenediamine tetraacetate, cooled to 0°C and mixed with 148 μl of 1,1,1-trifluoroacetone as well as a mixture that consists of 22 mg of oxone and 41 mg of sodium bicarbonate. It is allowed to react for 5 hours, poured onto sodium thiosulfate solution and extracted several times with ethyl acetate. The combined organic extracts are washed with saturated sodium chloride solution, and the residue that is obtained after filtration and removal of the solvent is purified by chromatography on an analytic thin-layer plate. As a mobile

55

5

10

solvent, a mixture of n-hexane and ethyl acetate is used. 3.2 mg
(6 μ mol, 39%) of title compound A and 1.0 mg (2 μ mol, 12%) of
title compound B are isolated in each case as a colorless oil.

15

$^1\text{H-NMR}$ (CDCl_3) of A: δ = 1.00 (3H), 1.02 (3H), 1.21-1.82
(7H), 1.29 (3H), 1.36 (3H), 1.95-2.06 (2H), 2.11 (3H), 2.30 (1H),
2.40 (1H), 2.48-2.62 (2H), 2.72 (3H), 2.81 (2H), 3.50 (1H), 3.69
(1H), 4.27 (1H), 4.52 (1H), 5.01 (1H), 5.06 (1H), 5.46 (1H), 5.72
(1H), 6.59 (1H), 6.99 (1H) ppm.

20

$^1\text{H-NMR}$ (CDCl_3) of B: δ = 0.96 (3H), 1.00 (3H), 1.20-1.91
(8H), 1.29 (3H), 1.34 (3H), 2.04 (1H), 2.09 (3H), 2.33 (1H),
2.42-2.61 (3H), 2.76 (3H), 2.93 (1H), 2.96 (1H), 3.38 (1H), 3.68
(1H), 3.99 (1H), 4.29 (1H), 4.98 (1H), 5.01 (1H), 5.57 (1H), 5.74
(1H), 6.69 (1H), 7.01 (1H) ppm.

25

30

Example 11

35

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-
methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-
en-1-yl)-cyclohexadec-13-ene-2,6-dione

40

Example 11a

45

(3S,6R,7S,8S,12E,15S,16E)-3,7-Bis-[[dimethyl(1,1-
dimethylethyl)silyl]oxy]-15-hydroxy-4,4,8,12,16-pentamethyl-17-

50

55

5

143

10

(2-methylthiazol-4-yl)-5-oxo-6-(prop-2-en-1-yl)-heptadeca-12,16-dienoic acid

15

Analogously to Example 1e, 230 mg (0.26 mmol) of compound B that is produced according to Example 9h is reacted, and after working-up, 214 mg (max. 0.26 mmol) of the title compound is isolated as a pale yellow oil, which is further reacted without

20

purification.

25

Example 11b

30

(4S,7R,8S,9S,13E,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

35

Analogously to Example 1g, 214 mg (max. 0.26 mmol) of the compound that is produced according to Example 11a is reacted, and after working-up and purification, 114 mg (0.15 mmol, 59%) of the title compound is isolated as a colorless oil.

40

¹H-NMR (CDCl₃): δ = 0.05 (3H), 0.08 (3H), 0.10 (3H), 0.13 (3H), 0.82-0.94 (21H), 1.12 (3H), 1.15-2.62 (13H), 1.21 (3H), 1.59 (3H), 2.11 (3H), 2.71 (3H), 3.03 (1H), 3.87 (1H), 4.30 (1H), 4.99 (1H), 5.03 (1H), 5.21 (1H), 5.28 (1H), 5.79 (1H), 6.51 (1H), 6.91 (1H) ppm.

45

50

55

5

144

10

Example 11c

15

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

20

Analogously to Example 1, 15 mg (20 μ mol) of the compound that is produced according to Example 11b is reacted, and after working-up and purification, 7.3 mg (14 μ mol, 71%) of the title compound is isolated as a colorless oil.

25

$^1\text{H-NMR}$ (CDCl_3): δ = 0.80-2.62 (13H), 0.99 (3H), 1.01 (3H), 1.26 (3H), 1.60 (3H), 2.04 (3H), 2.69 (3H), 3.49 (1H), 3.73 (1H), 4.01 (1H), 4.12 (1H), 4.42 (1H), 4.95-5.10 (3H), 5.37 (1H), 5.71 (1H), 6.56 (1H), 6.99 (1H) ppm.

30

Example 12

35

(1R,3S(E),7S,10R,11S,12S,16R)-7,11-Dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A)
and (1S,3S(E),7S,10R,11S,12S,16S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

45

50

Analogously to Example 10, 7.3 mg (14 μ mol) of the compound that is produced according to Example 11 is reacted, and after working-up and purification, 2.3 mg (4.3 μ mol, 31%) of title

55

5

145

10

compound A (or B) and 2.0 mg (3.7 μ mol, 27%) of title compound B (or A) are isolated in each case as a colorless oil.

15

$^1\text{H-NMR}$ (CDCl_3) of A (or B): δ = 0.90-2.34 (10H), 0.95 (3H), 1.01 (3H), 1.29 (3H), 1.38 (3H), 2.10 (3H), 2.47-2.62 (3H), 2.72 (3H), 2.88 (2H), 3.48 (1H), 3.80 (1H), 4.19 (1H), 4.32 (1H), 5.02 (1H), 5.07 (1H), 5.48 (1H), 5.77 (1H), 6.63 (1H), 7.00 (1H) ppm.

20

$^1\text{H-NMR}$ (CDCl_3) of B (or A): δ = 0.97 (3H), 1.06 (3H), 1.20-2.12 (9H), 1.25 (3H), 1.34 (3H), 2.08 (3H), 2.28 (1H), 2.46-2.62 (3H), 2.72 (3H), 2.92 (2H), 3.40 (1H), 3.68 (1H), 3.75 (1H), 4.28 (1H), 5.01 (1H), 5.06 (1H), 5.44 (1H), 5.72 (1H), 6.62 (1H), 6.99 (1H) ppm.

25

30

Example 13

35

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

40

Example 13a

45

(3RS,4S)-4-(2-Methyl-3-hydroxy-8-(trimethylsilyl)-hept-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

50

Analogously to Example 1a, 7.0 g (37 mmol) of (4S)-4-(2-methyl-1-oxo-prop-2-yl)-2,2-dimethyl-[1,3]dioxane, which was produced analogously to the process described in DE 197 51 200.3,

55

is reacted with 4-trimethylsilyl-but-3-in-1-yl-magnesium bromide, and after working-up and purification, 4.9 g (15.7 mmol, 42%) of the title compound is isolated as a colorless oil.

Example 13b

(4S)-4-(2-Methyl-3-oxo-8-(trimethylsilyl)-hept-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane

Analogously to Example 1b, 4.87 g (15.6 mmol) of the compound that is produced according to Example 13a is reacted, and after working-up and purification, 4.10 g (13.2 mmol, 85%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.13 (9H), 1.08 (3H), 1.13 (3H), 1.32 (1H), 1.34 (3H), 1.41 (3H), 1.61 (1H), 2.45 (2H), 2.73 (2H), 3.84 (1H), 3.96 (1H), 4.02 (1H) ppm.

Example 13c

(4S(4R,5S,6S,10E/Z,13S,14E))-4-(13-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-tetramethyl-3-oxo-15-(2-pyridyl)-4-(4-(trimethylsilyl)-prop-2-in-1-yl)-pentadec-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and
(4S(4S,5R,6S,10E/Z,13S,14E))-4-(13-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-5-hydroxy-2,6,10,14-tetramethyl-3-oxo-

5

10

15-(2-pyridyl)-4-(4-(trimethylsilyl)-prop-2-in-1-yl)-pentadec-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

15

20

25

Analogously to Example 1c, 2.74 g (8.82 mmol) of the compound that is produced according to Example 13b is reacted with 3.02 g (7.27 mmol) of (2S,6E/Z,9S,10E)-2,6,10-trimethyl-9-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-1-oxo-11-(2-pyridyl)-undeca-6,10-diene, which was produced analogously to the process that is described in DE 197 51 200.3, and after working-up and purification, in addition to 50% starting material, 1.63 g (2.2 mmol, 31%) of title compound A and 0.50 g (0.69 mmol, 9%) of title compound B are isolated in each case as a colorless oil.

30

35

¹H-NMR (CDCl₃) of A: δ = 0.00-0.20 (15H), 0.83-0.95 (12H), 1.00-1.80 (20H), 1.60+1.68 (3H), 1.90-2.10 (1H), 2.05 (3H), 2.28 (2H), 2.41 (1H), 2.55 (1H), 3.03+3.09 (1H), 3.46 (1H), 3.52 (1H), 3.78-4.20 (4H), 5.18 (1H), 6.49 (1H), 7.09 (1H), 7.23 (1H), 7.63 (1H), 8.60 (1H) ppm.

40

45

¹H-NMR (CDCl₃) of B: δ = 0.00-0.20 (15H), 0.86-1.00 (12H), 1.00-1.76 (19H), 1.61+1.70 (3H), 1.90-2.10 (2H), 2.06 (3H), 2.29 (2H), 2.53 (2H), 3.04 (1H), 3.43 (1H), 3.61 (1H), 3.80-4.18 (4H), 5.18 (1H), 6.48 (1H), 7.09 (1H), 7.23 (1H), 7.62 (1H), 8.59 (1H) ppm.

50

55

5

Example 13d

10

(3S,6R,7S,8S,12E/Z,15S,16E)-15-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-1,3,7-trihydroxy-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dien-5-one

15

Analogously to Example 1k, 2.25 g (3.10 mmol) of the compound that is produced according to Example 13c is reacted, and after working-up and purification, in addition to starting material, 1.31 g (1.91 mmol, 62%) of the title compound is isolated as a colorless oil.

20

25

¹H-NMR (CDCl₃): δ = 0.00-0.19 (9H), 0.85-0.98 (12H), 1.03-2.43 (25H), 1.60+1.69 (3H), 2.00+2.02 (3H), 2.69 (1H), 3.01+3.10 (1H), 3.31-3.60 (3H), 3.84 (2H), 4.02-4.26 (2H), 5.10+5.26 (1H), 6.41 (1H), 7.13 (1H), 7.32 (1H), 7.68 (1H), 8.61 (1H) ppm.

30

35

Example 13e

40

(3S,6R,7S,8S,12E/Z,15S,16E)-1,3,7,15-Tetrakis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dien-5-one

45

Analogously to Example 1l, 1.49 g (2.17 mmol) of the compound that is produced according to Example 13d is reacted,

50

55

5

10

and after working-up and purification, 1.95 g (1.90 mmol, 87%) of the title compound is isolated as a colorless oil.

15

20

¹H-NMR (CDCl₃): δ = 0.00-0.18 (33H), 0.86-0.98 (39H), 1.01-1.73 (7H), 1.08 (3H), 1.26 (3H), 1.61+1.69 (3H), 1.90-2.09 (2H), 2.05 (3H), 2.29 (2H), 2.51 (2H), 3.29 (1H), 3.53-3.71 (2H), 3.79 (1H), 3.89 (1H), 4.11 (1H), 5.17 (1H), 6.48 (1H), 7.09 (1H), 7.23 (1H), 7.61 (1H), 8.60 (1H) ppm.

25

Example 13f

30

(3S,6R,7S,8S,12E/Z,15S,16E)-1-Hydroxy-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dien-5-one

35

Analogously to Example 1m, 1.95 g (1.89 mmol) of the compound that is produced according to Example 13e is reacted, and after working-up and purification, 1.56 g (1.71 mmol, 90%) of the title compound is isolated as a colorless oil.

40

45

¹H-NMR (CDCl₃): δ = 0.00-0.17 (27H), 0.86-0.99 (30H), 1.07-1.78 (8H), 1.11 (3H), 1.26 (3H), 1.60+1.69 (3H), 1.90-2.09 (2H), 2.04 (3H), 2.29 (2H), 2.48 (1H), 2.68 (1H), 3.27 (1H), 3.66 (2H), 3.80 (1H), 4.11 (2H), 5.18 (1H), 6.49 (1H), 7.09 (1H), 7.22 (1H), 7.62 (1H), 8.60 (1H) ppm.

50

55

5

Example 13g

10

(3S,6R,7S,8S,12E/Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dienal

15

Analogously to Example 1n, 1.56 g (1.71 mmol) of the compound that is produced according to Example 13f is reacted, and after working-up, 1.61 g (max. 1.71 mmol) of the title compound is isolated as a yellow oil, which is further reacted without purification.

25

Example 13h

30

(3S,6R,7S,8S,12Z,15S,16E)-3,7,15-Tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dienoic acid (A) and (3S,6R,7S,8S,12E,15S,16E)-3,7,15-tris-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-heptadeca-12,16-dienoic acid (B)

35

40

The solution of 1.51 g (max. 1.60 mmol) of the compound, produced according to Example 13g, in 57 ml of tert-butanol is mixed with 47 ml of 2-methyl-2-butene, cooled to 2°C, mixed with 12.9 ml of water, 685 mg of sodium dihydrogen phosphate, 1.16 g

45

50

55

5
10 of sodium chlorite, allowed to heat to 23°C and stirred for 3
hours. It is poured into saturated sodium thiosulfate solution,
diluted with water and extracted several times with ethyl
15 acetate. The combined organic extracts are dried on sodium
sulfate, and the residue that is obtained after filtration and
removal of the solvent is purified by chromatography on fine
20 silica gel with a gradient system that consists of n-hexane and
ethyl acetate. 749 mg (807 μ mol, 50%) of title compound A and
579 mg (623 μ mol, 39%) of title compound B are isolated in each
25 case as a colorless oil.

¹H-NMR (CDCl₃) of A: δ = -0.02-0.17 (27H), 0.76-1.72 (6H),
0.88 (27H), 0.94 (3H), 1.10 (3H), 1.29 (3H), 1.68 (3H), 1.91-2.60
30 (7H), 2.02 (3H), 2.91 (1H), 3.39 (1H), 3.81 (1H), 4.11 (1H), 4.31
(1H), 5.18 (1H), 6.51 (1H), 7.09 (1H), 7.23 (1H), 7.62 (1H), 8.60
(1H) ppm.

35 ¹H-NMR (CDCl₃) of B: δ = 0.00-0.17 (27H), 0.80-0.98 (30H),
0.98-1.68 (6H), 1.08 (3H), 1.30 (3H), 1.60 (3H), 1.83-2.85 (8H),
40 2.05 (3H), 3.39 (1H), 3.79 (1H), 4.11 (1H), 4.30 (1H), 5.18 (1H),
6.48 (1H), 7.08 (1H), 8.22 (1H), 7.62 (1H), 8.60 (1H) ppm.

5

Example 13i

10

(3S,6R,7S,8S,12Z,15S,16E)-15-Hydroxy-3,7-bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(prop-2-in-1-yl)-heptadeca-12,16-dienoic acid

15

Analogously to Example 1e, 726 mg (782 μ mol) of compound A that is produced according to Example 13h is reacted, and after working-up, 657 mg (max. 782 μ mol) of the title compound is isolated, which is further reacted without purification.

20

25

Example 13j

(4S,7R,8S,9S,13Z,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

30

Analogously to Example 1q, 657 mg (max. 782 μ mol) of the compound that is produced according to Example 13i is reacted, and after working-up and purification, 300 mg (414 μ mol, 53%) of the title compound is isolated as a colorless oil.

35

40

$^1\text{H-NMR}$ (CDCl_3): δ = -0.08 (3H), 0.10 (3H), 0.15 (3H), 0.19 (3H), 0.81-2.20 (8H), 0.86 (9H), 0.95 (9H), 1.02 (3H), 1.14 (3H), 1.23 (3H), 1.68 (3H), 2.14 (3H), 2.33-2.82 (6H), 3.12 (1H), 4.06 (1H), 4.11 (1H), 5.02 (1H), 5.19 (1H), 6.58 (1H), 7.11 (1H), 7.26 (1H), 7.63 (1H), 8.59 (1H) ppm.

45

50

55

Example 13k

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 1, 140 mg (193 μ mol) of the compound that is produced according to Example 13j is reacted, and after working-up and purification, 52 mg (105 μ mol, 54%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 1.08 (3H), 1.10 (3H), 1.20-1.92 (6H), 1.42 (3H), 1.68 (3H), 2.02 (1H), 2.08 (3H), 2.22-2.72 (7H), 2.86 (1H), 3.43 (1H), 3.78 (1H), 4.37 (1H), 4.54 (1H), 5.12 (1H), 5.20 (1H), 6.61 (1H), 7.13 (1H), 7.30 (1H), 7.69 (1H), 8.55 (1H) ppm.

Example 14

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

Example 14a

(3S,6R,7S,8S,12E,15S,16E)-15-Hydroxy-3,7-bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4,4,8,12,16-pentamethyl-5-oxo-17-(2-pyridyl)-6-(prop-2-in-1-yl)-heptadeca-12,16-dienoic acid

5

154

10

Analogously to Example 1e, 534 mg (575 μ mol) of compound B that is produced according to Example 13h is reacted, and after working-up, 434 mg (max. 585 μ mol) of the title compound is isolated, which is further reacted without purification.

15

Example 14b

20

(4S,7R,8S,9S,13E,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)-silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

25

Analogously to Example 1q, 434 mg (max. 585 μ mol) of the compound that is produced according to Example 14a is reacted, and after working-up and purification, 382 mg (527 μ mol, 90%) of the title compound is isolated as a colorless oil.

30

$^1\text{H-NMR}$ (CDCl_3): δ = 0.04 (3H), 0.07-0.12 (9H), 0.85 (9H), 0.88 (9H), 0.93 (3H), 1.00-2.20 (8H), 1.14 (3H), 1.22 (3H), 1.58 (3H), 2.00 (1H), 2.12 (3H), 2.44-2.62 (5H), 3.19 (1H), 3.91 (1H), 4.41 (1H), 5.19 (1H), 5.29 (1H), 6.53 (1H), 7.09 (1H), 7.18 (1H), 7.62 (1H), 8.59 (1H) ppm.

40

Example 14c

45

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

50

55

Analogously to Example 1, 110 mg (152 μ mol) of the compound that is produced according to Example 14b is reacted, and after working-up and purification, 48 mg (97 μ mol, 64%) of the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.89-1.80 (5H), 1.01 (3H), 1.06 (3H), 1.35 (3H), 1.61 (3H), 1.93 (1H), 2.00 (1H), 2.10 (3H), 2.17 (1H), 2.38-2.66 (6H), 3.58 (1H), 3.79 (2H), 3.88 (1H), 4.44 (1H), 5.10 (1H), 5.40 (1H), 6.59 (1H), 7.13 (1H), 7.33 (1H), 7.68 (1H), 8.56 (1H) ppm.

Example 15

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Example 15a

(4S,7R,8S,9S,13Z,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione (A) and (4S,7R,8S,9S,13Z,16S(RS))-4,8-bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione (B)

5

156

10

15

20

25

The solution of 150 mg (207 μ mol) of the compound, produced according to Example 13j, in 16 ml of ethyl acetate is mixed with a catalytic amount of palladium on barium sulfate, 153 μ l of pyridine, and it is hydrogenated at 23°C under an atmosphere of hydrogen. After filtration and removal of the solvent, the residue is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. In addition to starting material, 66 mg (91 μ mol, 44%) of title compound A and 64 mg (88 μ mol, 42%) of title compound B are isolated in each case as an isolated oil.

30

35

$^1\text{H-NMR}$ (CDCl_3) of A: δ = -0.09 (3H), 0.07 (3H), 0.11 (6H), 0.78-1.82 (7H), 0.84 (9H), 0.92 (9H), 0.98 (3H), 1.09 (3H), 1.18 (3H), 1.67 (3H), 2.06-2.82 (7H), 2.13 (3H), 3.11 (1H), 4.02 (1H), 4.85-5.03 (3H), 5.18 (1H), 5.78 (1H), 6.57 (1H), 7.09 (1H), 7.25 (1H), 7.62 (1H), 8.59 (1H) ppm.

40

45

Example 15b

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

50

55

Analogously to Example 1, 65.6 mg (90 μ mol) of compound A that is produced according to Example 15a is reacted, and after

5

10

working-up and purification, 24.6 mg (49 μ mol, 55%) of the title compound is isolated as a colorless oil.

15

20

$^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (6H), 1.19-1.89 (5H), 1.32 (3H), 1.69 (3H), 2.05 (3H), 2.13-2.57 (6H), 2.64 (1H), 2.82 (1H), 3.33 (1H), 3.71 (2H), 4.34 (1H), 4.62 (1H), 5.01 (1H), 5.05 (1H), 5.12 (1H), 5.19 (1H), 5.75 (1H), 6.60 (1H), 7.12 (1H), 7.29 (1H), 7.68 (1H), 8.52 (1H) ppm.

Example 16

25

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

30

Example 16a

35

(4S,7R,8S,9S,13E,16S(E))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

40

45

Analogously to Example 15a, 114 g (157 μ mol) of the compound that is produced according to Example 14b is reacted, and after working-up and purification, 68 mg (94 μ mol, 60%) of the title compound is isolated as a colorless oil.

50

55

5

10

15

¹H-NMR (CDCl₃): δ = 0.04 (3H), 0.08 (3H), 0.10 (3H), 0.13 (3H), 0.83-0.98 (24H), 1.11 (3H), 1.15-1.96 (6H), 1.20 (3H), 2.08-2.65 (7H), 2.14 (3H), 3.03 (1H), 3.88 (1H), 4.31 (1H), 4.98 (1H), 5.02 (1H), 5.22 (1H), 5.29 (1H), 5.79 (1H), 6.54 (1H), 7.09 (1H), 7.20 (1H), 7.62 (1H), 8.60 (1H) ppm.

20

Example 16b

25

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

30

Analogously to Example 1, 67.7 mg (93 μmol) of the compound that is produced according to Example 16a is reacted, and after working-up and purification, 36.8 mg (74 μmol, 80%) of the title compound is isolated as a colorless oil.

35

¹H-NMR (CDCl₃): δ = 0.96-2.66 (13H), 0.99 (6H), 1.28 (3H), 1.62 (3H), 2.10 (3H), 3.49 (1H), 3.72 (1H), 4.01 (2H), 4.43 (1H), 4.91-5.13 (3H), 5.39 (1H), 5.71 (1H), 6.58 (1H), 7.12 (1H), 7.34 (1H), 7.66 (1H), 8.53 (1H) ppm.

40

Example 17

45

(1S/1R,3S(E),7S,10R(RS),11S,12S,16R/S)-7,11-Dihydroxy-10-(2,3-epoxyprop-1-yl)-3-(1-methyl-2-(2-N-oxido-pyridyl)ethenyl)-

50

55

5

10

8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

15

20

Analogously to Example 10, 36 mg (74 μ mol) of the compound that is produced according to Example 16 is reacted, and after working-up and purification, 12 mg (22 μ mol, 30%) of a mixture of two diastereomers A and B and 20 mg (37 μ mol, 50%) of a mixture of two diastereomers C and D of the title compounds are isolated in each case as a colorless oil.

25

MS (FAB): m/e = 546 ($M^+ + 1$)

Example 18

30

35

40

(1S,3S(E),7S,10R(R or S),11S,12S,16R)-7,11-Dihydroxy-10-(2,3-epoxyprop-1-yl)-3-(1-methyl-2-(2-N-oxido-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and (1R,3S(E),7S,10R(R or S),11S,12S,16S)-7,11-dihydroxy-10-(2,3-epoxyprop-1-yl)-3-(1-methyl-2-(2-N-oxido-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

45

50

The solution of 20 mg (37 μ mol) of a mixture of compounds C and D, produced according to Example 17, in 3.1 ml of anhydrous trichloromethane is mixed with a molecular sieve (4A), 789 ml of isopropanol, 14.2 mg of tetrapropylammonium perruthenate, and it is stirred for 5 hours at 55°C under an atmosphere of dry argon.

55

5

10

15

20

It is concentrated by evaporation, and the crude product that is obtained is purified by chromatography on analytical thin-layer plates. As a mobile solvent, a mixture of ethanol and ethyl acetate is used; as an eluant, a mixture of dichloromethane and ethanol is used. 4.6 mg (8.7 μ mol, 23%) of title compound A or B and 3.3 mg (6.2 μ mol, 17%) of title compound B or A are isolated in each case as a colorless oil.

25

$^1\text{H-NMR}$ (CDCl_3) of A or B: δ = 0.96 (3H), 1.06 (3H), 1.12-2.03 (11H), 1.22 (3H), 1.30 (3H), 2.11 (3H), 2.22 (1H), 2.58 (2H), 2.76 (1H), 3.44 (1H), 3.52 (1H), 3.73-3.91 (2H), 4.08-4.21 (2H), 4.47 (1H), 5.59 (1H), 6.59 (1H), 7.11 (1H), 7.23 (1H), 7.63 (1H), 8.59 (1H) ppm.

30

35

$^1\text{H-NMR}$ (CDCl_3) of B or A: δ = 0.96 (3H), 1.05 (3H), 1.11-1.96 (9H), 1.23 (3H), 1.31 (3H), 2.12 (3H), 2.19-2.35 (3H), 2.50-2.66 (2H), 2.78 (1H), 3.50-3.69 (3H), 3.93 (1H), 4.16 (1H), 4.25 (1H), 4.41 (1H), 5.59 (1H), 6.60 (1H), 7.12 (1H), 7.22 (1H), 7.64 (1H), 8.59 (1H) ppm.

40

Example 19

45

(1S/R,3S(E),7S,10R(S or R),11S,12S,16R/S)-7,11-Dihydroxy-10-(2,3-epoxyprop-1-yl)-3-(1-methyl-2-(2-N-oxido-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

50

55

5

10

15

20

25

30

35

45

50

55

Analogously to Example 18, 6.3 mg (12 μ mol) of compounds A and B that are produced according to Example 17 is reacted, and after working-up and purification, 2.4 mg (4.5 μ mol, 38%) of a mixture of the title compounds is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.95-2.22 (11H), 1.01 (3H), 1.10 (3H), 1.27 (3H), 1.31 (3H), 2.11 (3H), 2.34 (1H), 2.45-2.57 (2H), 2.90 (1H), 3.39-3.87 (4H), 4.01-4.37 (3H), 5.49 (1H), 6.62 (1H), 7.13 (1H), 7.24 (1H), 7.66 (1H), 8.58 (1H) ppm.

Example 20

(4S,7R,8S,9S,13Z,16S(R or S))-4,8-Dihydroxy-16-(1-methyl-2-(2-pyridyl)ethyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione (A) and (4S,7R,8S,9S,13Z,16S(S or R))-4,8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione (B)

Analogously to Example 1, 7.0 mg (9.6 μ mol) of compounds B that are produced according to Example 15a is reacted, and after working-up and purification, 1.4 mg (2.8 μ mol, 29%) of title compound A and 1.7 mg (3.4 μ mol, 35%) of title compound B are isolated in each case as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) of A: δ = 0.88 (1H), 0.92 (3H), 1.04 (3H), 1.07 (3H), 1.18-2.57 (14H), 1.30 (3H), 1.68 (3H), 2.91 (1H), 3.17

(1H), 3.28 (1H), 3.68 (1H), 4.47 (1H), 4.91-5.10 (4H), 5.70 (1H),
7.13-7.22 (2H), 7.68 (2H), 8.46 (1H) ppm.

¹H-NMR (CDCl₃) of B: δ = 1.00 (6H), 1.05 (3H), 1.10-2.59
(15H), 1.33 (3H), 1.63 (3H), 2.93 (1H), 3.11 (1H), 3.28 (1H),
3.63 (1H), 4.44 (1H), 4.91-5.12 (4H), 5.79 (1H), 6.39 (1H), 7.18
(2H), 7.67 (1H), 8.46 (1H) ppm.

Example 21

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(2-methyl-benzoxazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Example 21a

(2E/Z)-3-(2-Methyl-benzoxazol-5-yl)-2-propenoic acid ethyl ester

The suspension of 58 g (346 mmol) of 5-chloro-2-methylbenzoxazole, 200 ml of dimethylformamide, 57 g of sodium iodide and 16.2 g of nickel(II) bromide is heated for 4 hours to 150°C. After cooling, it is mixed with 42 ml of acrylic acid ethyl ester, 53 ml of triethylamine, 998 mg of tris-(dibenzylidene acetone)-dipalladium (0), 36.4 g of triphenylphosphine, and it is heated for three days to 150°C. The cooled mixture is poured into water, acidified and extracted several times with ethyl acetate. The combined organic extracts

5

10

15

are washed with saturated sodium chloride solution, dried on sodium sulfate, and the residue that is obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 6.4 g (28 mmol, 8%) of the title compound is isolated as a crystalline solid.

20

$^1\text{H-NMR}$ (CDCl_3): δ = 1.33 (3H), 2.64 (3H), 4.28 (2H), 6.42 (1H), 7.47 (2H), 7.78 (1H), 7.81 (1H) ppm.

25

Example 21b

(2-Methylbenzoxazol-5-yl)-carbaldehyde

30

35

40

45

50

The solution of 9.5 g (41 mmol) of the compound, produced according to Example 21a, in ml of tetrahydrofuran, is mixed with ml of water, ml of a 2.5% solution of osmium tetroxide in tert-butanol, g of sodium periodate, and it is stirred for 6 hours at 23°C. It is poured onto saturated sodium thiosulfate solution and extracted several times with ethyl acetate. The combined organic extracts are washed with saturated sodium chloride solution, dried on sodium sulfate, and the residue that is obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 4.86 g (30 mmol, 74%) of the title compound is isolated as a crystalline solid.

55

¹H-NMR (CDCl₃): δ = 2.69 (3H), 7.60 (1H), 7.90 (1H), 8.16 (1H), 10.08 (1H) ppm.

Example 21c

(3RS)-3-(2-Methyl-benzoxazol-5-yl)-1-[(4S,5R)-4-methyl-5-phenyl-oxazolidin-2-on-3-yl]-3-hydroxypropyl-1-one

50 ml of a 2.4 molar solution of n-butyllithium in n-hexane is added in drops at -30°C under an atmosphere of dry argon to the solution of 14.1 ml of diisopropylamine in 670 ml of anhydrous tetrahydrofuran, it is stirred for 20 minutes, cooled to -70°C and mixed within 4.5 hours with the solution of 19.8 g (4S,5R)-3-acetyl-4-methyl-5-phenyloxazolidin-2-one in 670 ml of tetrahydrofuran. After 1 hour, the solution of 4.86 g (30.1 mmol) of the compound, produced according to Example 21b, in 175 ml of tetrahydrofuran is added in drops within 1.5 hours, and it is stirred for 1 hour at -70°C. It is poured onto a saturated ammonium chloride solution, extracted several times with ethyl acetate, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate.

11.3 g (29.7 mmol, 98%) of the title compound is isolated as a colorless oil.

Example 21d

(3S)-3-(2-Methyl-benzoxazol-5-yl)-1-[(4S,5R)-4-methyl-5-phenyl-oxazolidin-2-on-3-yl]-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-propyl-1-one (A) and (3R)-3-(2-methyl-benzoxazol-5-yl)-1-[(4S,5R)-4-methyl-5-phenyl-oxazolidin-2-on-3-yl]-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-propyl-1-one (B)

The solution of 12.5 g (32.8 mmol) of the compound, produced according to Example 21c, in 110 ml of anhydrous dichloromethane is cooled under an atmosphere of dry argon to -70°C, mixed with 7.8 ml of 2,6-lutidine and 13.9 ml of trifluoromethanesulfonic acid-tert-butyldimethylsilylester, and it is stirred for 1 hour. It is poured onto a saturated sodium bicarbonate solution, extracted several times with dichloromethane, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is separated by chromatography on fine silica gel with a gradient system that consists of n-hexane, ethyl acetate and ethanol. 8.9 g (18.0 mmol, 55%) of title compound A is isolated as a crystalline

5

10

solid, and 2.9 g (5.9 mmol, 18%) of title compound B is isolated as a colorless oil.

15

¹H-NMR (CDCl₃) of A: δ = -0.19 (3H), 0.02 (3H), 0.82 (9H), 0.88 (3H), 2.61 (3H), 3.19 (1H), 3.51 (1H), 4.69 (1H), 5.36 (1H), 5.55 (1H), 7.21-7.44 (7H), 7.64 (1H) ppm.

20

¹H-NMR (CDCl₃) of B: δ = -0.19 (3H), 0.04 (3H), 0.85 (9H), 0.88 (3H), 2.63 (3H), 3.04 (1H), 4.67 (1H), 4.77 (1H), 5.39 (1H), 5.63 (1H), 7.21-7.46 (7H), 7.67 (1H) ppm.

25

Example 21e

30

(3S)-3-(2-Methyl-benzoxazol-5-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-propionic acid ethyl ester

35

The solution of 13.9 g (28.2 mmol) of the compound, produced according to Example 21d, in 140 ml of anhydrous ethanol is mixed at 23°C under an atmosphere of dry argon with 7.1 ml of titanium tetraethylate, and it is heated for 3 hours to 85°C. It is concentrated by evaporation, and the residue is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 10.1 g (27.8 mmol, 99%) of the title compound is isolated as a colorless oil.

45

¹H-NMR (CDCl₃): δ = -0.20 (3H), 0.02 (3H), 0.82 (9H), 1.26 (3H), 2.55 (1H), 2.62 (3H), 2.76 (1H), 4.12 (2H), 5.26 (1H), 7.29 (1H), 7.40 (1H), 7.62 (1H) ppm.

50

55

5

Example 21f

10

(3S)-3-(2-Methyl-benzoxazol-5-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-propan-1-ol

15

The solution of 10.1 g (27.8 mmol) of the compound, produced according to Example 21e, in ml of anhydrous dichloromethane, is cooled under an atmosphere of dry argon to -78°C, mixed with 58 ml of a 1.2 molar solution of diisobutylaluminum hydride in toluene, and it is stirred for 1 more hour. It is mixed with 16 ml of isopropanol, 32 ml of water, allowed to heat to 23°C and stirred until a fine-grained precipitate has formed. After filtration and removal of the solvent, 7.2 g (22.4 mmol, 81%) of the title compound is isolated as a colorless oil.

25

30

¹H-NMR (CDCl₃): δ = -0.18 (3H), 0.07 (3H), 0.89 (9H), 1.97 (2H), 2.35 (1H), 2.66 (3H), 3.73 (2H), 5.06 (1H), 7.28 (1H), 7.42 (1H), 7.60 (1H) ppm.

35

Example 21g

40

(3S)-3-(2-Methyl-benzoxazol-5-yl)-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-1-iodo-propane

45

The solution of 2.83 g of triphenylphosphine in 40 ml of anhydrous dichloromethane is mixed at 23°C under an atmosphere of dry argon with 737 mg of imidazole, 2.71 g of iodine, and the solution of 2.65 g (8.2 mmol) of the compound, produced according

50

55

5

10

15

20

25

30

35

40

45

50

55

to Example 21f, in 30 ml of dichloromethane, is added in drops while being cooled. It is stirred for 1 hour and purified directly by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 2.3 g (5.3 mmol, 65%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = -0.20 (3H), 0.06 (3H), 0.85 (9H), 2.10 (1H), 2.21 (1H), 2.61 (3H), 3.11 (1H), 3.23 (1H), 4.82 (1H), 7.22 (1H), 7.39 (1H), 7.59 (1H) ppm.

Example 21h

(3S)-3-(2-Methyl-benzoxazol-5-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-propane-1-triphenylphosphonium iodide

2.3 g (5.3 mmol) of the compound that is produced according to Example 21g is mixed with 2.9 ml of ethyldiisopropylamine, 17.5 g of triphenylphosphine, and it is heated for 4 hours to 85°C. The oily residue is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 3.3 g (4.8 mmol, 89%) of the title compound is isolated as a crystalline solid.

¹H-NMR (CDCl₃): δ = -0.19 (3H), 0.12 (3H), 0.84 (9H), 1.89 (1H), 2.09 (1H), 2.60 (3H), 3.41 (1H), 4.06 (1H), 5.37 (1H), 7.38 (1H), 7.49 (1H), 7.59 (1H), 7.62-7.84 (15H) ppm.

5

Example 21i

10

(2S,6E/Z,9S)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-9-(2-methyl-benzoxazol-5-yl)-1-(tetrahydropyran-2-yloxy)-2,6-dimethyl-non-6-ene

15

20

The solution of 2.3 g (3.3 mmol) of the compound, produced according to Example 21h, in 15 ml of anhydrous tetrahydrofuran is mixed under an atmosphere of dry argon at 0°C with 5 ml of a 1.0 molar solution of sodium hexamethyldisilazane in tetrahydrofuran, the solution of 513 mg (2.25 mmol) of (2S)-2-methyl-6-oxo-heptane-1-(tetrahydropyran-2-yloxy), which was produced analogously to the process described in DE 197 51 200.3, is added in drops in 15 ml of tetrahydrofuran, allowed to heat to 23°C and reacted for 3 more hours. It is poured onto a saturated ammonium chloride solution, extracted several times with ethyl acetate, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is separated by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 506 mg (1.0 mmol, 44%) of the title compound is isolated as a colorless oil.

35

40

45

50

¹H-NMR (CDCl₃): δ = -0.15 (3H), 0.01 (3H), 0.80-0.92 (12H), 1.02 (1H), 1.19-1.97 (12H), 1.46+1.62 (3H), 2.21-2.48 (2H), 2.60

55

(3H), 3.10+3.19 (1H), 3.40-3.61 (2H), 3.82 (1H), 4.53 (1H), 4.69 (1H), 5.11 (1H), 7.22 (1H), 7.37 (1H), 7.57 (1H) ppm.

Example 21j

(2S,6E/Z,9S)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-9-(2-methyl-benzoxazol-5-yl)-1-hydroxy-2,6-dimethyl-non-6-ene

Analogously to Example 1k, 447 mg (0.87 mmol) of the compound that is produced according to Example 21i is reacted, and after working-up and purification, 298 mg (0.69 mmol, 79%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = -0.12 (3H), 0.01 (3H), 0.82-0.92 (12H), 1.01 (1H), 1.16-1.67 (4H), 1.44+1.63 (3H), 1.83-1.98 (2H), 2.18 (1H), 2.33 (1H), 2.44 (1H), 2.62 (3H), 3.31-3.53 (2H), 4.71 (1H), 5.07+5.13 (1H), 7.24+7.29 (1H), 7.39 (1H), 7.53+7.58 (1H) ppm.

Example 21k

(2S,6E/Z,9S)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-9-(2-methyl-benzoxazol-5-yl)-1-oxo-2,6-dimethyl-non-6-ene

Analogously to Example 1n, 272 mg (0.63 mmol) of the compound that is produced according to Example 21j is reacted, and after working-up and purification, 236 mg (0.55 mmol, 87%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = -0.16 (3H), 0.01 (3H), 0.84 (9H),
1.02+1.05 (3H), 1.13-2.50 (9H), 1.44+1.61 (3H), 2.61 (3H), 4.71
(1H), 5.13 (1H), 7.21 (1H), 7.37 (1H), 7.55 (1H), 9.54 (1H) ppm.

Example 21l

(4S(4R,5S,6S,10E/Z,13S))-4-(13-[(1,1-Dimethylethyl)dimethylsilyl]oxy)-4-(prop-2-en-1-yl)-13-(2-methyl-benzoxazol-5-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-tridec-10-en-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and (4S(4S,5R,6S,10E/Z,13S))-4-(13-[(1,1-dimethylethyl)dimethylsilyl]oxy)-4-(prop-2-en-1-yl)-13-(2-methyl-benzoxazol-5-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-tridec-10-en-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

Analogously to Example 1c, 236 mg (0.55 mmol) of the compound that is produced according to Example 21k is reacted with 433 mg (1.80 mmol) of (4S)-4-(2-methyl-3-oxo-hept-6-en-2-yl)-2,2-dimethyl-[1,3]dioxane, which was produced analogously to the process that is described in DE 197 51 200.3, and after working-up and purification, in addition to starting material, 221 mg (0.33 mmol, 60%) of title compound A and 72 mg (0.11 mmol, 20%) of title compound B are isolated in each case as a colorless oil.

¹H-NMR (CDCl₃) of A: δ = -0.13 (3H), 0.01 (3H), 0.78-0.88 (12H), 0.96 (3H), 1.04 (1H), 1.11-2.52 (12H), 1.23 (3H), 1.31

5

10

15

(3H), 1.39 (3H), 1.47+1.64 (3H), 2.62 (3H), 2.90+2.98 (1H), 3.32 (1H), 3.47 (1H), 3.87 (1H), 3.97 (1H), 4.13 (1H), 4.70 (1H), 4.98 (1H), 5.03 (1H), 5.12 (1H), 5.71 (1H), 7.22 (1H), 7.38 (1H), 7.56 (1H) ppm.

Example 21m

20

(3S,6R,7S,8S,12E/Z,15S)-15-[[[(1,1-Dimethylethyl)dimethylsilyl]oxy]-6-(prop-2-en-1-yl)-1,3,7-trihydroxy-4,4,8,12-tetramethyl-15-(2-methyl-benzoxazol-5-yl)-pentadec-12-en-5-one

25

Analogously to Example 1k, 221 mg (0.33 mmol) of compound A that is produced according to Example 21l is reacted, and after working-up and purification, 163 mg (0.26 mmol, 78%) of the title compound is isolated as a colorless oil.

30

35

¹H-NMR (CDCl₃): δ = -0.15 (3H), 0.01 (3H), 0.79-0.90 (12H), 1.05 (3H), 1.17-2.59 (13H), 1.20+1.24 (3H), 1.43+1.62 (3H), 2.62+2.64 (3H), 2.81+3.07 (1H), 3.25-3.70 (3H), 3.86 (2H), 4.08 (2H), 4.68 (1H), 4.92-5.19 (3H), 5.69 (1H), 7.25+7.29 (1H), 7.39 (1H), 7.48+7.52 (1H) ppm.

40

45

Example 21n

50

55

5

10

(3S,6R,7S,8S,12E/Z,15S)-6-(Prop-2-en-1-yl)-1,3,7,15-tetrakis-
[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-15-
(2-methyl-benzoxazol-5-yl)-pentadec-12-en-5-one

15

Analogously to Example 1l, 163 mg (0.26 mmol) of the
compound that is produced according to Example 2lm is reacted,
and after working-up and purification, 236 mg (0.24 mmol, 93%) of
the title compound is isolated as a colorless oil.

20

25

¹H-NMR (CDCl₃): δ = -0.06 (3H), -0.04-0.08 (21H), 0.79-0.93
(39H), 0.96-1.66 (7H), 1.01 (3H), 1.17 (3H), 1.47+1.62 (3H), 1.88
(2H), 2.18-2.52 (4H), 2.61 (3H), 3.11 (1H), 3.53 (1H), 3.63 (1H),
3.73 (1H), 3.84 (1H), 4.68 (1H), 4.91 (1H), 4.97 (1H), 5.12 (1H),
5.72 (1H), 7.21 (1H), 7.36 (1H), 7.56 (1H) ppm.

30

Example 21o

35

(3S,6R,7S,8S,12E/Z,15S)-1-Hydroxy-6-(prop-2-en-1-yl)-3,7,15-tris-
[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-15-
(2-methyl-benzoxazol-5-yl)-pentadec-12-en-5-one

40

Analogously to Example 1m, 236 mg (0.24 mmol) of the
compound that is produced according to Example 2ln is reacted,
and after working-up and purification, 146 mg (0.17 mmol, 71%) of
the title compound is isolated as a colorless oil.

45

50

55

5

Example 21p

10

(3S,6R,7S,8S,12E/Z,15S)-5-Oxo-6-(prop-2-en-1-yl)-3,7,15-tris-
[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-15-
(2-methyl-benzoxazol-5-yl)-pentadec-12-enal

15

Analogously to Example 1n, 146 mg (0.17 mmol) of the
compound that is produced according to Example 21o is reacted,
and after working-up and purification, 143 mg (0.17 mmol, 98%) of
the title compound is isolated as a colorless oil.

20

25

Example 21q

(3S,6R,7S,8S,12Z,15S)-5-Oxo-6-(prop-2-en-1-yl)-3,7,15-tris-
[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-5-
(2-methyl-benzoxazol-5-yl)-pentadec-12-enoic acid (A) and
(3S,6R,7S,8S,12E,15S)-5-oxo-6-(prop-2-en-1-yl)-3,7,15-tris-
[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl)-15-
(2-methyl-benzoxazol-5-yl)-pentadec-12-enoic acid (B)

30

35

The solution of 143 mg (0.17 mmol) of the compound, produced
according to Example 21p, in 5 ml of tert-butanol is mixed at 0°C
with the solution of 1.1 ml of 2-methyl-2-butene in 3.6 ml of
tetrahydrofuran, 1.3 ml of water, 67 mg of sodium dihydrogen
phosphate, 117 mg of sodium chlorite, and it is stirred for 2
hours. It is poured onto a saturated sodium thiosulfate
solution, extracted several times with ethyl acetate, the

40

45

50

55

5

10

15

20

combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is separated by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate. 58 mg (66 μ mol, 39%) of title compound A and 52 mg (60 μ mol, 35%) of title compound B are isolated in each case as a colorless oil.

25

$^1\text{H-NMR}$ (CDCl_3) of A: δ = -0.13 (3H), -0.02 (6H), 0.04 (6H), 0.12 (3H), 0.80-0.92 (27H), 0.96 (3H), 1.06 (3H), 1.09-1.96 (7H), 1.15 (3H), 1.70 (3H), 2.13-2.60 (7H), 2.62 (3H), 3.20 (1H), 3.66 (1H), 4.43 (1H), 4.72 (1H), 4.92 (1H), 4.99 (1H), 5.26 (1H), 5.70 (1H), 7.34 (1H), 7.40 (1H), 7.89 (1H) ppm.

30

$^1\text{H-NMR}$ (CDCl_3) of B: δ = -0.11 (3H), 0.02 (6H), 0.07 (3H), 0.10 (3H), 0.16 (3H), 0.86-0.94 (30H), 0.90-2.05 (8H), 1.12 (3H), 1.19 (3H), 1.39 (3H), 2.23-2.60 (6H), 2.63 (3H), 3.21 (1H), 3.79 (1H), 4.36 (1H), 4.68 (1H), 4.98 (1H), 5.01 (1H), 5.10 (1H), 5.77 (1H), 7.36 (1H), 7.41 (1H), 7.54 (1H) ppm.

35

40

Example 21r

45

(3S,6R,7S,8S,12Z,15S)-15-Hydroxy-5-oxo-6-(prop-2-en-1-yl)-3,7-bis-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-15-(2-methyl-benzoxazol-5-yl)-pentadec-12-enoic acid

50

55

5

10

15

Analogously to Example 1p, 58 mg (66 μ mol) of compound A that is produced according to Example 21q is reacted, and after working-up, 52 mg (max. 66 μ mol) of the title compound is isolated, which is further reacted without purification.

Example 21s

20

(4S,7R,8S,9S,13Z,16S)-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(2-methyl-benzoxazol-5-yl)-7-(prop-2-en-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

25

Analogously to Example 1q, 52 mg (max. 66 μ mol) of the compound that is produced according to Example 21r is reacted, and after working-up and purification, 42 mg (57 μ mol, 86%) of the title compound is isolated as a colorless oil.

30

35

$^1\text{H-NMR}$ (CDCl_3): δ = -0.08 (3H), 0.09 (6H), 0.14 (3H), 0.77-1.88 (7H), 0.85 (9H), 0.93 (9H), 1.01 (3H), 1.09 (3H), 1.15 (3H), 1.71 (3H), 2.10-2.75 (6H), 2.62 (3H), 2.91 (1H), 3.11 (1H), 4.00 (1H), 4.92 (1H), 4.99 (1H), 5.19 (1H), 5.57 (1H), 5.79 (1H), 7.32 (1H), 7.44 (1H), 7.68 (1H) ppm.

40

Example 21t

45

(4S,7R,8S,9S,13Z,16S)-4,8-Dihydroxy-16-(2-methyl-benzoxazol-5-yl)-7-(prop-2-en-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

50

55

5

10

Analogously to Example 1, 42 mg (57 μ mol) of the compound that is produced according to Example 21s is reacted, and after working-up and purification, 19 mg (37 μ mol, 65%) of the title compound is isolated as a colorless oil.

15

20

$^1\text{H-NMR}$ (CDCl_3): δ = 1.02 (3H), 1.08 (3H), 1.14-1.97 (6H), 1.22 (3H), 1.70 (3H), 2.22-2.60 (7H), 2.62 (3H), 2.78-2.95 (2H), 3.36 (1H), 3.78 (1H), 4.10 (1H), 5.03 (1H), 5.09 (1H), 5.19 (1H), 5.76 (1H), 5.85 (1H), 7.28 (1H), 7.43 (1H), 7.63 (1H) ppm.

25

Example 22

30

(4S,7R,8S,9S,13E,16S)-4,8-Dihydroxy-16-(2-methyl-benzoxazol-5-yl)-7-(prop-2-en-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

35

Example 22a

40

(3S,6R,7S,8S,12E,15S)-15-Hydroxy-5-oxo-6-(prop-2-en-1-yl)-3,7-bis-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4,4,8,12-tetramethyl-15-(2-methyl-benzoxazol-5-yl)-pentadec-12-enoic acid

45

Analogously to Example 1p, 52 mg (60 μ mol) of compound B that is produced according to Example 21q is reacted, and after working-up, 46 mg (max. 60 μ mol) of the title compound is isolated, which is further reacted without purification.

50

55

5

Example 22b

10

(4S,7R,8S,9S,13E,16S)-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(2-methyl-benzoxazol-5-yl)-7-(prop-2-en-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

15

Analogously to Example 1q, 46 mg (max. 60 μ mol) of the compound that is produced according to Example 22a is reacted, and after working-up and purification, 32 mg (43 μ mol, 72%) of the title compound is isolated as a colorless oil.

20

$^1\text{H-NMR}$ (CDCl_3): δ = 0.03-0.11 (12H), 0.89 (9H), 0.91 (9H), 0.94-1.96 (6H), 0.98 (3H), 1.12 (3H), 1.21 (3H), 1.59 (3H), 2.10-2.76 (7H), 2.63 (3H), 3.08 (1H), 3.91 (1H), 4.31 (1H), 5.02 (1H), 5.07 (1H), 5.29 (1H), 5.79 (1H), 5.89 (1H), 7.30 (1H), 7.42 (1H), 7.62 (1H) ppm.

25

30

Example 22c

35

(4S,7R,8S,9S,13E,16S)-4,8-Dihydroxy-16-(2-methyl-benzoxazol-5-yl)-7-(prop-2-en-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

40

Analogously to Example 1, 32 mg (43 μ mol) of the compound that is produced according to Example 22b is reacted, and after working-up and purification, 15 mg (29 μ mol, 68%) of the title compound is isolated as a colorless oil.

45

50

55

5

10

15

¹H-NMR (CDCl₃): δ = 0.99 (3H), 1.02 (3H), 1.27 (3H), 1.38-1.99 (6H), 1.64 (3H), 2.18 (1H), 2.23-2.76 (6H), 2.62 (3H), 3.34 (1H), 3.49 (2H), 3.75 (1H), 4.32 (1H), 4.96-5.08 (3H), 5.73 (1H), 5.98 (1H), 7.23 (1H), 7.42 (1H), 7.67 (1H) ppm.

Example 23

20

(4S,7R,8S,9S,13Z,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

25

Example 23a

30

(4S(4R,5S,6S,10E/Z,13S,14Z))-4-(13-[[[(1,1-Dimethylethyl)dimethylsilyl]oxy]-4-(prop-2-in-1-yl)-14-fluoro-15-(2-methylthiazol-4-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-pentadeca-(10,14-dien-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and
(4S(4S,5R,6S,10E/Z,13S,14Z))-4-(13-[[[(1,1-Dimethylethyl)dimethylsilyl]oxy]-4-(prop-2-in-1-yl)-14-fluoro-15-(2-methylthiazol-4-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-pentadeca-10,14-dien-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

40

45

Analogously to Example 1c, 2.89 g (6.57 mmol) of (2S,6E/Z,9S,10Z)-9-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-10-fluoro-11-(2-methyl-4-thiazolyl)-2,6-dimethylundeca-6,10-dienal, which was produced analogously to the process that is described

50

55

in DE 19907480.1, is reacted with 5.09 g (16.4 mmol) of (4S)-4-(2-methyl-3-oxo-7-trimethylsilyl-hept-6-in-2-yl)-2,2-dimethyl-[1,3]dioxane, which was produced according to the process described in DE 19751200.3, and after working-up and purification, in addition to starting material, 3.26 g (4.35 μ mol, 66%) of title compound A as well as 602 mg (0.80 mmol, 12%) of title compound B are isolated in each case as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) of A: δ = 0.03-0.13 (15H), 0.82-0.92 (12H), 0.97-2.08 (12H), 1.06 (3H), 1.30 (6H), 1.38 (3H), 1.58+1.65 (3H), 2.33-2.47 (3H), 2.55 (1H), 2.70 (3H), 3.44 (1H), 3.52 (1H), 3.80-4.28 (2H), 5.13 (1H), 6.03 (1H), 7.32 (1H) ppm.

$^1\text{H-NMR}$ (CDCl_3) of B: δ = 0.05-0.65 (15H), 0.88-0.99 (12H), 1.02-1.73 (8H), 1.18 (6H), 1.32 (3H), 1.41 (3H), 1.60+1.69 (3H), 1.90-2.08 (2H), 2.33-2.58 (4H), 2.70 (3H), 3.43 (1H), 3.60 (1H), 3.79-4.26 (4H), 5.18 (1H), 6.05 (1H), 7.33 (1H) ppm.

Example 23b

(3S,6R,7S,8S,12E/Z,15S,16Z)-15-[[[(1,1-Dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-16-fluoro-1,3,7-trihydroxy-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-12,16-dien-5-one

Analogously to Example 1k, 3.26 g (4.35 mmol) of compound A that is produced according to Example 23a is reacted, and after

working-up and purification, in addition to starting material,
2.44 g (3.43 μ mol, 79%) of the title compound is isolated as a
colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.03-0.15 (15H), 0.85-0.95 (12H), 0.98-
2.08 (8H), 1.14 (3H), 1.26 (3H), 1.58+1.67 (3H), 2.31-2.49 (3H),
2.59-2.76 (2H), 2.72 (3H), 2.89 (1H), 3.06 (1H), 3.42 (1H), 3.47
3.58 (2H), 3.88 (2H), 4.08-4.22 (2H), 5.11+5.18 (1H), 5.98 (1H),
7.33 (1H) ppm.

Example 23c

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-1,3,7,15-tetrakis-[[[1,1-
dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-
1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-
12,16-dien-5-one

Analogously to Example 11, 2.77 g (3.90 mmol) of the
compound that is produced according to Example 23b is reacted,
and after working-up and purification, 3.48 g (3.31 mmol, 85%) of
the title compound is isolated as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.00-0.15 (33H), 0.83-0.97 (39H), 1.00-
1.75 (7H), 1.07 (3H), 1.27 (3H), 1.60+1.68 (3H), 1.88-2.03 (2H),
2.31-2.48 (2H), 2.51 (2H), 2.70 (3H), 3.29 (1H), 3.52-3.71 (2H),
3.29 (1H), 3.89 (1H), 4.19 (1H), 5.15 (1H), 6.06 (1H), 7.33 (1H)
ppm.

5

Example 23d

10

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-1-hydroxy-3,7,15-tris-

[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-

15

prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-
heptadeca-12,16-dien-5-one

Analogously to Example 1m, 3.48 g (3.31 mmol) of the

20

compound that is produced according to Example 23c is reacted,
and after working-up and purification, 2.36 g (2.5 mmol, 76%) of
the title compound is isolated as a colorless oil.

25

 $^1\text{H-NMR}$ (CDCl_3): δ = 0.00-0.18 (27H), 0.83-0.99 (30H), 1.01-
1.80 (7H), 1.12 (3H), 1.27 (3H), 1.60+1.68 (3H), 1.86-2.07 (3H),
2.83-2.52 (3H), 2.64 (1H), 2.70 (3H), 3.26 (1H), 3.66 (2H), 3.80
30 (1H), 4.10 (1H), 4.20 (1H), 5.16 (1H), 6.06 (1H), 7.32 (1H) ppm.

Example 23e

35

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-5-oxo-3,7,15-tris-[[(1,1-
dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-
40 1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-
12,16-dienal

Analogously to Example 1n, 2.36 g (2.51 mmol) of the

45

compound that is produced according to Example 23d is reacted,
and after working-up and purification, 2.25 g (2.40 mmol, 96%) of
the title compound is isolated as a colorless oil.

50

55

5

Example 23f

10

(3S,6R,7S,8S,12Z,15S,16Z)-16-Fluoro-5-oxo-3,7,15-tris-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-12,16-dienoic acid (A) and

15

20

(3S,6R,7S,8S,12E,15S,16Z)-16-fluoro-5-oxo-3,7,15-tris-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-1,2,16-dienoic acid (B)

25

Analogously to Example 22q, 2.25 g (2.40 mmol) of the compound that is produced according to Example 23e is reacted, and after working-up and purification, 960 mg (1.01 mmol, 42%) of title compound A as well as 937 mg (0.98 mmol, 41%) of title compound are isolated in each case as a colorless oil.

30

35

¹H-NMR (CDCl₃) of A: δ = -0.02-0.17 (27H), 0.89 (27H), 0.94 (3H), 1.08-1.67 (6H), 1.18 (3H), 1.22 (3H), 1.70 (3H), 1.89 (1H), 2.12 (1H), 2.28-2.53 (5H), 2.61 (1H), 2.69 (3H), 3.31 (1H), 3.71 (1H), 4.20 (1H), 4.38 (1H), 5.18 (1H), 6.40 (1H), 7.36 (1H) ppm.

40

45

¹H-NMR (CDCl₃) of B: δ = -0.01-0.18 (27H), 0.84-0.97 (30H), 1.00-1.55 (6H), 1.20 (3H), 1.23 (3H), 1.59 (3H), 1.82-2.05 (2H), 2.25-2.60 (4H), 2.65 (1H), 2.70 (3H), 3.33 (1H), 3.76 (1H), 4.16 (1H), 4.38 (1H), 5.13 (1H), 6.12 (1H), 7.38 (1H) ppm.

50

55

5

Example 23g

10

(3S,6R,7S,8S,12Z,15S,16Z)-16-Fluoro-5-oxo-3,7-bis-[[(1,1-dimethylethyl)dimethylsilyl]oxy]-15-hydroxy-6-(prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-12,16-dienoic acid

15

Analogously to Example 1e, 960 mg (1.01 mmol) of compound A that is produced according to Example 23f is reacted, and after working-up, 898 mg (max. 1.01 mmol) of the title compound is isolated, which is further reacted without purification.

20

25

Example 23h

30

(4S,7R,8S,9S,13Z,16S(Z))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-fluoro-2-(2-methylthiazol-4-yl)-ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

35

Analogously to Example 1q, a total of 896 mg (max. 1.01 mmol) of the compound that is produced according to Example 23b is reacted in several portions, and after working-up and purification, 480 mg (0.64 mmol, 64%) of the title compound is isolated as a colorless oil.

40

45

¹H-NMR (CDCl₃): δ = -0.10 (3H), 0.12 (3H), 0.15 (3H), 0.19 (3H), 0.80-1.83 (96H), 0.85 (9H), 0.94 (9H), 1.01 (3H), 1.18 (3H),

50

55

5

10

1.23 (3H), 1.68 (3H), 2.08 (1H), 2.22-2.89 (7H), 2.69 (3H), 3.09 (1H), 4.00-4.12 (2H), 5.07-5.21 (2H), 6.13 (1H), 7.36 (1H) ppm.

15

Example 23i

20

(4S,7R,8S,9S,13Z,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

25

Analogously to Example 1, 60 mg (80 μ mol) of the compound that is produced according to Example 23h is reacted, and after working-up and purification, 28 mg (54 μ mol, 67%) of the title compound is isolated as a colorless oil.

30

$^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (3H), 1.11 (3H), 1.18-1.42 (3H), 1.38 (3H), 1.56-1.97 (3H), 1.90 (3H), 2.05 (1H), 2.28 (1H), 2.33-2.66 (6H), 2.69 (3H), 2.79 (1H), 3.30 (1H), 3.38 (1H), 3.79 (1H), 4.21 (1H), 5.12 (1H), 5.46 (1H), 6.19 (1H), 7.36 (1H) ppm.

35

Example 24

40

(4S,7R,8S,9S,13E,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-12,6-dione

45

50

55

5

Example 24a

10

(3S,6R,7S,8S,12E,15S,16Z)-16-Fluoro-5-oxo-3,7-bis-[[[1,1-dimethylethyl)dimethylsilyl]oxy]-15-hydroxy-6-(prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methylthiazol-4-yl)-heptadeca-12,16-dienoic acid

15

Analogously to Example 1e, 937 mg (0.98 mmol) of compound B that is produced according to Example 23f is reacted, and after working-up, 914 mg (max. 0.98 mmol) of the title compound is isolated, which is further reacted without purification.

25

Example 24b

30

(4S,7R,8S,9S,13E,16S(Z))-4,8-Bis-[[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-fluoro-2-(2-methylthiazol-4-yl)-ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

35

Analogously to Example 1q, 914 mg (max. 0.98 mmol) of the compound that is produced according to Example 24a is reacted, and after working-up and purification, 451 mg (603 μ mol, 62%) of the title compound is isolated as a colorless oil.

40

$^1\text{H-NMR}$ (CDCl_3): δ = 0.02-0.12 (12H), 0.79-1.73 (5H), 0.89 (18H), 0.96 (3H), 1.12 (3H), 1.22 (3H), 1.58 (3H), 1.91 (1H), 2.01 (1H), 2.11 (1H), 2.39-2.80 (6H), 2.69 (3H), 3.15 (1H), 3.91 (1H), 4.33 (1H), 5.17 (1H), 5.42 (1H), 6.12 (1H), 7.36 (1H) ppm.

50

55

5

Example 24c

10

(4S,7R,8S,9S,13E,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

15

Analogously to Example 1, 451 mg (603 μ mol) of the compound that is produced according to Example 24b is reacted, and after working-up and purification, 170 mg (327 μ mol, 54%) of the title compound is isolated as a colorless oil.

20

$^1\text{H-NMR}$ (CDCl_3): δ = 0.86 (1H), 1.00 (3H), 1.03 (3H), 1.26-2.23 (7H), 1.33 (3H), 1.60 (3H), 2.41-2.62 (6H), 2.69 (3H), 3.59 (1H), 3.79 (1H), 4.02-4.19 (2H), 4.39 (1H), 5.11 (1H), 5.54 (1H), 6.17 (1H), 7.37 (1H) ppm.

25

30

Example 25

35

(1S,3S(Z),7S,10R,11S,12S,16R)-7,11-Dihydroxy-3-(1-fluoro-2-(2-methyl-4-thiazolyl)ethenyl)-10-(prop-2-in-1-yl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A)
and (1R,3S(Z),7S,10R,11S,12S,16S)-7,11-dihydroxy-3-(1-fluoro-2-(2-methyl-4-thiazolyl)ethenyl)-10-(prop-2-in-1-yl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

40

45

The solution of 50 mg (96 μ mol) of the compound, produced according to Example 23, in 4.5 ml of acetonitrile is mixed at 0°C with 554 μ l of a 0.1 M aqueous solution of ethylenediamine

50

55

5
10 tetraacetate, 638 μ l of trifluoroacetone, 260 mg of sodium
bicarbonate, 150 mg of oxone, and it is stirred for 1.5 hours at
23°C. It is mixed with sodium thiosulfate solution, extracted
15 several times with ethyl acetate, the combined organic extracts
are washed with saturated sodium chloride solution, dried on
sodium sulfate, and the residue is purified by chromatography on
20 analytic thin-layer plates. As a mobile solvent, a mixture that
consists of dichloromethane and isopropanol is used; as an
eluant, a mixture that consists of dichloromethane and methanol
25 is used. 29 mg (54 μ mol, 56%) of title compound A as well as
9 mg (17 μ mol, 18%) of title compound B are isolated in each case
as a colorless oil.

30 $^1\text{H-NMR}$ (CDCl_3) of A: δ = 1.01 (3H), 1.08 (3H), 1.22-1.81
(7H), 1.28 (3H), 1.39 (3H), 2.01 (1H), 2.04 (1H), 2.19 (1H),
35 2.40-2.76 (5H), 2.69 (3H), 2.91 (1H), 3.60 (1H), 3.80 (1H), 4.19
(1H), 4.31 (1H), 5.70 (1H), 6.23 (1H), 7.38 (1H) ppm.

40 $^1\text{H-NMR}$ (CDCl_3) of B: δ = 0.97 (3H), 1.08 (3H), 1.19-1.96
(8H), 1.25 (3H), 1.42 (3H), 1.99 (1H), 2.28 (1H), 2.42-2.62 (4H),
2.70 (3H), 2.98 (1H), 3.04 (1H), 3.49 (1H), 3.62 (1H), 4.04 (1H),
45 4.23 (1H), 5.80 (1H), 6.21 (1H), 7.38 (1H) ppm.

Example 26

(1S,3S(Z),7S,10R,11S,12S,16S)-7,11-Dihydroxy-3-(1-fluoro-2-(2-methyl-4-thiazolyl)ethenyl)-10-(prop-2-in-1-yl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A)
and (1R,3S(Z),7S,10R,11S,12S,16R)-7,11-dihydroxy-3-(1-fluoro-2-(2-methyl-4-thiazolyl)ethenyl)-10-(prop-2-in-1-yl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)
and (1SR,3S(Z),7S,10R,11S,12S,16SR)-7,11-dihydroxy-3-(1-fluoro-2-(2-methyl-4-(N-oxido)-thiazolyl)ethenyl)-10-(prop-2-in-1-yl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (C).

Analogously to Example 25, 80 mg (154 μmol) of the compound that is produced according to Example 24 is reacted, and after working-up and purification, 21 mg (39 μmol , 25%) of title compound A, 31 mg (58 μmol , 38%) of title compound B as well as 3 mg (6 μmol , 4%) of title compounds C are isolated in each case as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) of A or B: δ = 0.96 (3H), 1.08 (3H), 1.18-1.85 (7H), 1.22 (3H), 1.38 (3H), 1.99 (1H), 2.09 (1H), 2.20 (1H), 2.40 (1H), 2.51-2.72 (3H), 2.68 (3H), 2.99 (1H), 3.13 (1H), 3.53 (1H), 3.75 (1H), 3.82 (1H), 4.30 (1H), 5.66 (1H), 6.21 (1H), 7.37 (1H) ppm.

5

10

15

¹H-NMR (CDCl₃) of B or A: δ = 0.93 (3H), 1.04 (3H), 1.11-1.81 (7H), 1.28 (3H), 1.41 (3H), 1.99 (1H), 2.06-2.23 (2H), 2.43 (1H), 2.51-2.72 (4H), 2.69 (3H), 2.87 (1H), 3.55 (1H), 3.85 (1H), 4.19 (1H), 4.31 (1H), 5.66 (1H), 6.24 (1H), 7.39 (1H) ppm.

20

¹H-NMR of C (CDCl₃): δ = 0.95+0.99 (3H), 1.08+1.10 (3H), 1.13-2.77 (14H), 1.22+1.26 (3H), 1.45+1.51 (3H), 2.59 (3H), 2.95 (1H), 3.52-3.86 (2H), 4.13+5.41 (1H), 4.43-4.70 (2H), 5.63+5.72 (1H), 6.56+6.59 (1H), 7.41+7.46 (1H) ppm.

25

Example 27

30

(4S,7R,8S,9S,13Z,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

35

Example 27a

4-(2-Methyloxazolyl)-carbaldehyde

40

The solution of 36.6 g (236 mmol) of 4-(2-methyloxazolyl)-carboxylic acid ethyl ester in 795 ml of anhydrous dichloromethane is cooled under an atmosphere of dry argon to -78°C, mixed with 378 ml of a 1.0 molar solution of diisobutylaluminum hydride in n-hexane, and it is stirred for 1 more hour. It is mixed with 96 ml of isopropanol, 160 ml of water, allowed to heat to 23°C and stirred until a fine-grained

50

55

5

10

precipitate has formed. After filtration and removal of the solvent, 24.7 g (222 mmol, 94%) of the title compound is isolated as a pale yellow oil.

15

$^1\text{H-NMR}$ (CDCl_3): δ = 2.53 (3H), 8.17 (1H), 9.90 (1H) ppm.

Example 27b

20

(2Z)-3-(2-Methyloxazol-4-yl)-2-fluoro-2-propenoic acid ethyl ester (A) and (2E)-3-(2-methyloxazol-4-yl)-2-fluoro-2-propenoic acid ethyl ester (B)

25

30

The solution of 106 g of 2-fluoro-2-phosphonoacetic acid triethyl ester in 224 ml of ethylene glycol dimethyl ether is added in drops under an atmosphere of dry argon at 0°C to 19.1 g of a 55% sodium hydride dispersion in 224 ml of anhydrous ethylene glycol dimethyl ether, and it is stirred for one more hour. Then, it is mixed with the solution of 26.4 g (238 mmol) of the compound, produced according to Example 27a, in 224 ml of ethylene glycol dimethyl ether, and it is allowed to heat within 1 hour to 23°C. It is poured onto a saturated ammonium chloride solution, extracted several times with ethyl acetate, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is purified by chromatography on fine silica gel with a gradient

35

40

45

50

55

5
10 system that consists of n-hexane and ethyl acetate. 24.8 g (125 mmol, 52%) of title compound A is isolated as a crystalline solid, and 12.5 g (63 mmol, 26%) of title compound B is isolated as a colorless oil.

15 ¹H-NMR (CDCl₃) of A: δ = 1.37 (3H), 2.49 (3H), 4.32 (2H), 6.91 (1H), 7.94 (1H) ppm.

20 ¹H-NMR (CDCl₃) of B: δ = 1.39 (3H), 2.47 (3H), 4.36 (2H), 6.75 (1H), 8.53 (1H) ppm.

25 Example 27c

(2Z)-3-(2-Methyloxazol-4-yl)-2-fluoro-2-propenoic acid ethyl ester

30 The solution of 24.4 g (123 mmol) of compound B, produced according to Example 27b, in 130 ml of anhydrous toluene is mixed with 5.3 ml of thiophenol, and it is stirred for 2 days under an atmosphere of dry argon at 23°C. It is poured into a 5% sodium hydroxide solution, extracted several times with ethyl acetate, 40 the combined organic extracts are washed with water, saturated sodium chloride solution and dried on magnesium sulfate. The residue that is obtained after filtration and removal of the 45 solvent is purified by chromatography on fine silica gel with a gradient system that consists of n-hexane and ethyl acetate.

50

55

19.5 g (98 mmol, 80%) of the title compound is isolated as a crystalline solid.

Example 27d

(2Z)-3-(2-Methyloxazol-4-yl)-2-fluoro-2-propenal

The solution of 26.2 g (131 mmol) of compound A, produced according to Example 27b or 3c, in 380 ml of anhydrous toluene is cooled under an atmosphere of dry argon to -78°C, mixed with 180 ml of a 1.2 M solution of diisobutylaluminum hydride in toluene, and it is stirred for 8 hours. It is mixed with water, extracted several times with ethyl acetate, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. After filtration and removal of the solvent, 20.1 g (128 mmol, 98%) of the title compound is isolated as a colorless oil, which is further reacted without purification.

¹H-NMR (CDCl₃): δ = 2.51 (3H), 6.69 (1H), 8.07 (1H), 9.32 (1H) ppm.

Example 27e

(3S,4Z)-5-(2-Methyloxazol-4-yl)-1-[(4S,5R)-4-methyl-5-phenyl-oxazolidin-2-on-3-yl]-3-hydroxy-4-fluoro-4-penten-1-one (A) and (3R,4Z)-5-(2-methyloxazol-4-yl)-1-[(4S,5R)-4-methyl-5-phenyl-oxazolidin-2-on-3-yl]-3-hydroxy-4-fluoro-4-penten-1-one (B)

136 ml of a 2.4 molar solution of n-butyllithium in n-hexane is added in drops at -30°C under an atmosphere of dry argon to the solution of 45.8 ml of diisopropylamine in 2 l of anhydrous tetrahydrofuran, and it is stirred for 20 minutes, cooled to -70°C and mixed within 4 hours with the solution of 64.2 g of (4S,5R)-3-acetyl-4-methyl-5-phenyloxazolidin-2-one in 1 l of tetrahydrofuran. After 1 hour, the solution of 15.1 g (97.6 mmol) of the compound, produced according to Example 27d, in 650 ml of tetrahydrofuran is added in drops within 2 hours, and it is stirred for 16 hours at -70°C. It is poured onto a saturated ammonium chloride solution, extracted several times with ethyl acetate, the combined organic extracts are washed with saturated sodium chloride solution and dried on sodium sulfate. The residue that is obtained after filtration and removal of the solvent is separated by repeated chromatography on fine silica gel with a gradient system that consists of n-hexane, ethyl acetate and ethanol. 19.9 g (53 mmol, 54%) of title compound A is isolated as a crystalline solid, and 8.2 g (22 mmol, 22%) of title compound B is isolated as a colorless foam.

¹H-NMR (CDCl₃) of A: δ = 0.92 (3H), 2.47 (3H), 3.33 (1H), 3.50 (1H), 3.70 (1H), 4.73-4.88 (2H), 5.71 (1H), 5.97 (1H), 7.26-7.48 (5H), 7.75 (1H) ppm.

195.

¹H-NMR (CDCl₃) of B: δ = 0.93 (3H), 2.48 (3H), 3.40 (2H),
4.73-4.90 (2H), 5.70 (1H), 5.98 (1H), 7.24-7.49 (5H), 7.76 (1H)
ppm.

Example 27f

(3S,4Z)-5-(2-Methyloxazol-4-yl)-1-[(4S,5R)-4-methyl-5-phenyl-
oxazolidin-2-on-3-yl]-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-
4-fluoro-4-penten-1-one

Analogously to Example 11, 16.2 g (43.5 mmol) of compound A
that is produced according to Example 27e is reacted, and after
working-up and purification, 15.9 g (32.5 mmol, 75%) of the title
compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.11 (6H), 0.88 (9H), 0.90 (3H), 2.46
(3H), 3.24 (1H), 3.52 (1H), 4.77 (1H), 4.89 (1H), 5.66 (1H), 5.83
(1H), 7.23-7.48 (5H), 7.74 (1H) ppm.

Example 27g

(3S,4Z)-5-(2-Methyloxazol-4-yl)-3-[[dimethyl(1,1-
dimethylethyl)silyl]oxy]-4-fluoro-4-pentenoic acid ethyl ester

Analogously to Example 22e, 15.6 g (32.6 mmol) of the
compound that is produced according to Example 27f is reacted,
and after working-up and purification, 11.4 g (32 mmol, 98%) of
the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.08 (6H), 0.88 (9H), 1.26 (3H), 2.43 (3H), 2.67 (2H), 4.13 (2H), 4.71 (1H), 5.80 (1H), 7.72 (1H) ppm.

Example 27h

(3S,4Z)-5-(2-Methyloxazol-4-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4-fluoro-4-penten-1-ol

Analogously to Example 22f, 11.4 g (31.9 mmol) of the compound that is produced according to Example 27g is reacted, and after working-up and purification, 9.16 g (29 mmol, 91%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.07 (3H), 0.10 (3H), 0.90 (9H), 1.94 (2H), 2.08 (1H), 2.43 (3H), 3.73 (1H), 3.80 (1H), 4.49 (1H), 5.80 (1H), 7.71 (1H) ppm.

Example 27i

(3S,4Z)-5-(2-Methyloxazol-4-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-1-iodo-4-fluoro-4-pentene

Analogously to Example 22g, 7.16 g (22.7 mmol) of the compound that is produced according to Example 27h is reacted, and after working-up and purification, 8.06 g (18.9 mmol, 83%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.09 (3H), 0.15 (3H), 0.91 (9H), 2.20 (2H), 2.46 (3H), 3.23 (2H), 4.33 (1H), 5.80 (1H), 7.73 (1H) ppm.

Example 27j

(3S,4Z)-5-(2-Methyloxazol-4-yl)-3-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-4-fluoro-4-pentene-1-triphenylphosphonium iodide

Analogously to Example 22h, 8.06 g (18.9 mmol) of the compound that is produced according to Example 27i is reacted, and after working-up and purification, 10.7 g (15.6 mmol, 82%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.10 (3H), 0.18 (3H), 0.87 (9H), 1.97 (1H), 2.10 (1H), 2.42 (3H), 3.48 (1H), 3.97 (1H), 4.86 (1H), 5.93 (1H), 7.63-7.88 (16H) ppm.

Example 27k

(2S,6E/Z,9S,10Z)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-10-fluoro-11-(2-methyloxazol-4-yl)-1-(tetrahydropyran-2-yloxy)-2,6-dimethyl-undeca-6,10-diene

Analogously to Example 22i, 3.20 g (14.0 mmol) of the compound that is produced according to Example 27j is reacted, and after working-up and purification, 3.53 g (6.9 mmol, 49%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.08 (6H), 0.84-0.97 (12H), 1.09 (1H), 1.22-2.04 (12H), 1.59+1.68 (3H), 2.30-2.49 (2H), 2.44 (3H), 3.06-

3.27 (1H), 3.42-3.62 (2H), 3.86 (1H), 4.19 (1H), 4.55 (1H), 5.12
(1H), 5.73 (1H), 7.71 (1H) ppm.

Example 27l

(2S,6E/Z,9S,10Z)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-10-fluoro-11-(2-methyloxazol-4-yl)-1-hydroxy-2,6-dimethyl-undeca-6,10-diene

Analogously to Example 1k, 3.48 g (6.83 mmol) of the compound that is produced according to Example 27k is reacted, and after working-up and purification, 2.28 g (5.36 mmol, 78%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.08 (6H), 0.83-0.94 (12H), 1.03 (1H), 1.21-1.70 (5H), 1.58+1.68 (3H), 1.91-2.05 (2H), 2.27-2.50 (2H), 2.44 (3H), 3.37-3.52 (2H), 4.19 (1H), 5.12 (1H), 5.72 (1H), 7.72 (1H) ppm.

Example 27m

(2S,6E/Z,9S,10Z)-9-[[Dimethyl(1,1-dimethylethyl)silyl]oxy]-10-fluoro-11-(2-methyloxazol-4-yl)-2,6-dimethyl-undeca-6,10-dienal

Analogously to Example 1n, 2.28 g (5.36 mmol) of the compound that is produced according to Example 27l is reacted, and after working-up and purification, 2.27 g (5.36 mmol, 100%) of the title compound is isolated as a colorless oil.

5

10

¹H-NMR (CDCl₃): δ = 0.06 (6H), 0.90 (9H), 1.03+1.08 (3H),
1.21-1.46 (4H), 1.57+1.66 (3H), 2.00 (2H), 2.21-2.42 (3H), 2.45
(3H), 4.19 (1H), 5.14 (1H), 5.73 (1H), 7.71 (1H), 9.59 (1H) ppm.

15

Example 27n

20

(4S(4R,5S,6S,10E/Z,13S,14Z))-4-(13-[(1,1-
Dimethylethyl)dimethylsilyl]oxy]-4-(prop-2-in-1-yl)-14-fluoro-15-
(2-methyloxazol-4-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-pentadeca-
10,14-dien-2-yl)-2,2-dimethyl-[1,3]dioxane (A) and

25

(4S(4S,5R,6S,10E/Z,13S,14Z))-4-(13-[(1,1-
dimethylethyl)dimethylsilyl]oxy]-4-(prop-2-in-1-yl)-14-fluoro-15-
(2-methyloxazol-4-yl)-3-oxo-5-hydroxy-2,6,10-trimethyl-pentadeca-
10,14-dien-2-yl)-2,2-dimethyl-[1,3]dioxane (B)

30

35

Analogously to Example 1c, 1.87 g (4.41 mmol) of the
compound, produced according to Example 27m, with (4S)-4-(2-
methyl-3-oxo-7-trimethylsilyl-hept-6-in-2-yl)-2,2-dimethyl-
[1,3]dioxane, which was produced according to the process
described in DE 19751200.3, is reacted, and after working-up and
purification, in addition to starting material, 1.37 g (1.87
mmol, 42%) of title compound A as well as 190 mg (0.26 mmol, 6%)
of title compound B are isolated in each case as a colorless oil.

45

50

¹H-NMR (CDCl₃) of A: δ = 0.02-0.16 (15H), 0.81-0.93 (12H),
0.97-1.78 (13H), 1.06 (3H), 1.39 (3H), 1.58+1.67 (3H), 1.91-2.08

55

(2H), 2.30-2.48 (3H), 2.44 (3H), 2.55 (1H), 3.03 (1H), 3.45 (1H),
3.52 (1H), 3.88 (1H), 3.99 (1H), 4.08-4.23 (2H), 5.12 (1H), 5.72
(1H), 7.72 (1H) ppm.

¹H-NMR (CDCl₃) of B: δ = 0.01-0.12 (15H), 0.82-1.73 (18H),
0.89 (9H), 1.17 (3H), 1.40 (3H), 1.58+1.67 (3H), 1.88-2.05 (2H),
2.28-2.57 (3H), 2.42 (3H), 3.41 (1H), 3.59 (1H), 3.79-4.05 (3H),
4.18 (1H), 5.11 (1H), 5.72 (1H), 7.70 (1H) ppm.

Example 27o

(3S,6R,7S,8S,12E/Z,15S,16Z)-15-[[[(1,1-Dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-16-fluoro-1,3,7-trihydroxy-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dien-5-one

Analogously to Example 1k, 2.16 g (2.94 mmol) of compound A that is produced according to Example 27n is reacted, and after working-up and purification, 1.47 g (2.12 mmol, 72%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.03 (6H), 0.15 (9H), 0.85-0.95 (12H), 0.98-1.80 (7H), 1.15 (3H), 1.27 (3H), 1.57+1.66 (3H), 1.90-1.08 (2H), 2.30-2.45 (3H), 2.49+2.51 (3H), 2.58-2.72 (2H), 2.90+3.03 (1H), 3.37-3.72 (3H), 3.88 (2H), 4.07-4.22 (2H), 5.11 (1H), 5.63+5.70 (1H), 7.71 (1H) ppm.

5

Example 27p

10

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-1,3,7,15-tetrakis-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dien-5-one

15

Analogously to Example 11, 1.47 g (2.12 mmol) of the compound that is produced according to Example 27o is reacted, and after working-up and purification, 2.13 g (2.05 mmol, 97%) of the title compound is isolated as a colorless oil.

20

25

¹H-NMR (CDCl₃): δ = 0.00-0.15 (33H), 0.83-0.98 (39H), 1.00-1.72 (7H), 1.07 (3H), 1.26 (3H), 1.59+1.67 (3H), 1.94 (2H), 2.27-2.43 (2H), 2.43 (3H), 2.51 (2H), 3.28 (1H), 3.52-3.71 (2H), 3.78 (1H), 3.88 (1H), 4.18 (1H), 5.12 (1H), 5.73 (1H), 7.71 (1H) ppm.

30

Example 27q

35

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-1-hydroxy-3,7,15-tris-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dien-5-one

40

Analogously to Example 1m, 2.13 g (2.05 mmol) of the compound that is produced according to Example 27p is reacted, and after working-up and purification, 1.47 g (1.60 mmol, 78%) of the title compound is isolated as a colorless oil.

45

50

55

5

10

15

$^1\text{H-NMR}$ (CDCl_3): δ = 0.00-0.15 (27H), 0.83-0.98 (30H), 1.02-1.77 (7H), 1.10 (3H), 1.27 (3H), 1.59+1.68 (3H), 1.89-2.07 (3H), 2.30-2.52 (3H), 2.45 (3H), 1.68 (1H), 3.27 (1H), 3.60-3.71 (2H), 3.79 (1H), 4.05-4.23 (2H), 5.13 (1H), 5.73 (1H), 7.70 (1H) ppm.

Example 27r

20

25

(3S,6R,7S,8S,12E/Z,15S,16Z)-16-Fluoro-5-oxo-3,7,15-tris-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-yn-1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dienal

30

Analogously to Example 1n, 1.47 g (1.60 mmol) of the compound that is produced according to Example 27q is reacted, and after working-up, 1.62 g (max. 1.60 mmol) of the title compound is isolated as a colorless oil.

35

40

$^1\text{H-NMR}$ (CDCl_3) of a purified sample: δ = -0.01-0.11 (27H), 0.83-0.98 (30H), 1.00-1.56 (5H), 1.11 (3H), 1.28 (3H), 1.59+1.68 (3H), 1.88-2.02 (2H), 2.29-2.50 (4H), 2.43 (3H), 2.58-2.71 (2H), 3.26 (1H), 3.78 (1H), 4.18 (1H), 4.50 (1H), 5.12 (1H), 5.73 (1H), 7.71 (1H), 9.77 (1H) ppm.

45

Example 27s

50

(3S,6R,7S,8S,12Z,15S,16Z)-16-Fluoro-5-oxo-3,7,15-tris-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-

55

5

10

15

20

25

30

35

40

45

50

55

1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-
12,16-dienoic acid (A) and
(3S,6R,7S,8S,12E,15S,16Z)-16-fluoro-5-oxo-3,7,15-tris-[[[(1,1-
dimethylethyl)dimethylsilyl]oxy]-6-(3-(trimethylsilyl)-prop-2-in-
1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-
12,16-dienoic acid (B)

Analogously to Example 22q, 1.60 g (max. 1.60 mmol) of the
compound that is produced according to Example 27r is reacted,
and after working-up and purification, 601 mg (642 μ mol, 40%) of
title compound A as well as 500 mg (534 μ mol, 33%) of title
compound B are isolated in each case as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) of A: δ = -0.04-0.19 (27H), 0.89 (27H), 0.96
(3H), 1.05-2.53 (13H), 1.19 (3H), 1.26 (3H), 1.69 (3H), 2.46
(3H), 2.63 (1H), 3.32 (1H), 3.71 (1H), 4.61 (1H), 4.39 (1H), 5.18
(1H), 6.08 (1H), 7.73 (1H) ppm.

$^1\text{H-NMR}$ (CDCl_3) of B: δ = -0.02-0.18 (27H), 0.90 (30H),
0.99-2.67 (14H), 1.21 (6H), 1.58 (3H), 2.46 (3H), 3.32 (1H), 3.74
(1H), 4.13 (1H), 4.36 (1H), 5.10 (1H), 5.79 (1H), 7.72 (1H) ppm.

Example 27t

(3S,6R,7S,8S,12Z,15S,16Z)-16-Fluoro-5-oxo-3,7-bis-[[[(1,1-
dimethylethyl)dimethylsilyl]oxy]-15-hydroxy-6-(prop-2-in-1-yl)-

5
10 4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dienoic acid

15 Analogously to Example 1e, 601 mg (642 μ mol) of compound A that is produced according to Example 27s is reacted, and after working-up, 657 mg (max. 642 μ mol) of the title compound is isolated as a crude product, which is further reacted without
20 purification.

Example 27u

25 (4S,7R,8S,9S,13Z,16S(Z))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-
30 cyclohexadec-13-ene-2,6-dione

35 Analogously to Example 1q, 657 mg (max. 642 μ mol) of the compound that is produced according to Example 27u is reacted, and after working-up and purification, 91 mg (124 μ mol, 19%) of the title compound is isolated as a colorless oil.

40 Example 27v

45 (4S,7R,8S,9S,13Z,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione
50
55

5

10

Analogously to Example 1, 91 mg (124 μ mol) of the compound that is produced according to Example 27u is reacted, and after working-up and purification, 45 mg (89 μ mol, 73%) of the title compound is isolated as a colorless oil.

15

20

$^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (3H), 1.10 (3H), 1.20-1.42 (4H), 1.37 (3H), 1.58-1.94 (2H), 1.69 (3H), 2.04 (1H), 2.20-2.84 (8H), 2.45 (3H), 3.20 (1H), 3.38 (1H), 3.78 (1H), 4.20 (1H), 5.11 (1H), 5.43 (1H), 5.90 (1H), 7.73 (1H) ppm.

25

Example 28

30

(4S,7R,8S,9S,13E,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

35

Example 28a

40

(3S,6R,7S,8S,12E,15S,16Z)-16-Fluoro-5-oxo-3,7-bis-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-15-hydroxy-6-(prop-2-in-1-yl)-4,4,8,12-tetramethyl-17-(2-methyloxazol-4-yl)-heptadeca-12,16-dienoic acid

45

Analogously to Example 1e, 500 mg (534 μ mol) of compound B that is produced according to Example 27f is reacted, and after working-up, 517 mg (max. 534 μ mol) of the title compound is

50

55

isolated as crude product, which is further reacted without purification.

Example 28b

(4S,7R,8S,9S,13E,16S(Z))-4,8-Bis-[[dimethyl(1,1-dimethylethyl)silyl]oxy]-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

Analogously to Example 1q, 517 mg (534 μ mol) of the compound that is produced according to Example 28a is reacted, and after working-up and purification, 128 mg (175 μ mol, 33%) of the title compound is isolated as a colorless oil.

Example 28c

(4S,7R,8S,9S,13E,16S(Z))-4,8-Dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-7-(prop-2-in-1-yl)-1-oxa-5,5,9,13-tetramethyl-cyclohexadec-13-ene-2,6-dione

Analogously to Example 1, 128 mg (175 μ mol) of the compound that is produced according to Example 28b is reacted, and after working-up and purification, 54 mg (107 μ mol, .61%) of the title compound is isolated as a colorless oil.

¹H-NMR (CDCl₃): δ = 0.89 (1H), 0.98 (3H), 1.02 (3H), 1.20-2.23 (7H), 1.33 (3H), 1.59 (3H), 2.40-2.61 (6H), 2.42 (3H), 3.57

(1H), 3.77 (1H), 3.82 (1H), 3.87 (1H), 4.33 (1H), 5.08 (1H), 5.53
(1H), 5.87 (1H), 7.72 (1H) ppm.

Example 29

(4S,7R,8S,9S,13Z,16S(E))-4,8-Dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 21, 41.4 mg of the title compound is obtained from 5-chloro-2-methylbenzothiazole as a colorless oil.

¹H-NMR (CDCl₃): δ = 1.04 (3H), 1.07 (3H), 1.24 (3H), 1.72 (3H), 1.3-1.8 (3H), 1.89 (1H), 2.26-2.63 (7H), 2.84 (3H), 2.90 (2H), 3.36 (1H), 3.78 (1H), 4.12 (1H), 5.05 (1H), 5.07 (1H), 5.19 (1H), 5.76 (1H), 5.88 (1H), 7.34 (1H), 7.80 (1H), 7.95 (1H) ppm.

Example 30

(4S,7R,8S,9S,13E,16S(E))-4,8-Dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

Analogously to Example 22, 108.2 mg of the title compound is obtained from 5-chloro-2-methylbenzothiazole as a colorless oil.

¹H-NMR (CDCl₃): δ = 1.00 (3H), 1.05 (3H), 1.24 (3H), 1.66 (3H), 1.5-1.97 (3H), 1.75-1.99 (2H), 2.09-2.58 (7H), 2.79 (1H), 2.83 (3H), 3.56 (1H), 3.80 (1H), 3.86 (1H), 4.08 (1H), 4.49 (1H),

4.93 (1H), 5.00 (1H), 5.01 (1H), 5.73 (1H), 6.03 (1H), 7.28 (1H),
7.77 (1H), 8.00 (1H) ppm.

Example 31

(1S,3S(Z),7S,10R,11S,12S,16R)-7,11-Dihydroxy-3-(2-methyl-
benzothiazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(Z),7S,10R,11S,12S,16S)-7,11-dihydroxy-3-(2-methyl-
benzothiazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 10, 13.6 mg of title compound A and
4.5 mg of title compound B are obtained from 25.0 mg of the title
compound that is produced in Example 29.

¹H-NMR (CDCl₃) of title compound A: δ = 0.98 (3H), 1.02
(3H), 1.23 (3H), 1.32 (3H), 1.2-1.8 (7H), 2.18 (2H), 2.27 (1H),
2.43-2.69 (4H), 2.84 (3H), 2.93 (1H), 3.60 (1H), 3.69 (1H), 4.21
(1H), 4.44 (1H), 5.02 (1H), 5.06 (1H), 5.72 (1H), 6.19 (1H), 7.36
(1H), 7.82 (1H), 7.94 (1H) ppm.

¹H-NMR (CDCl₃) of title compound B: δ = 0.98 (3H), 1.00
(3H), 1.31 (3H), 1.34 (3H), 1.1-1.75 (6H), 1.83 (1H), 2.0-2.65
(6H), 2.84 (3H), 3.03 (1H), 3.06 (1H), 3.28-3.43 (2H), 4.03 (1H),
4.31 (1H), 4.98 (1H), 5.03 (1H), 5.75 (1H), 6.27 (1H), 7.36 (1H),
7.81 (1H), 7.97 (1H) ppm.

Example 32

(1S,3S(Z),7S,10R,11S,12S,16S)-7,11-Dihydroxy-3-(2-methyl-
benzothiazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(Z),7S,10R,11S,12S,16R)-7,11-dihydroxy-3-(2-methyl-
benzothiazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-
4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 10, 17.7 mg of title compound A and 14.6 mg of title compound B are obtained from 60.0 mg of the title compound that is produced in Example 30 by plate cleaning with a mixture that consists of methylene chloride/ethyl acetate at a 6:4 ratio.

¹H-NMR (CDCl₃) of title compound A: δ = 0.96 (3H), 1.01 (3H), 1.31 (3H), 1.38 (3H), 1.2-1.9 (7H), 2.01-2.15 (1H), 2.21-2.35 (3H), 2.46-2.65 (3H), 2.83 (3H), 2.93 (1H), 3.47 (1H), 3.83 (1H), 4.20-4.34 (2H), 5.02 (1H), 5.07 (1H), 5.79 (1H), 6.13 (1H), 7.36 (1H), 7.81 (1H), 7.96 (1H) ppm.

¹H-NMR (CDCl₃) of title compound B: δ = 1.01 (3H), 1.04 (3H), 1.14 (3H), 1.33 (3H), 1.1-1.75 (6H), 2.05-2.37 (4H), 2.42-2.65 (3H), 2.84 (3H), 2.88 (1H), 3.03 (1H), 3.42 (1H), 3.49 (1H), 3.79 (1H), 4.26 (1H), 5.02 (1H), 5.06 (1H), 5.74 (1H), 6.12 (1H), 7.32 (1H), 7.80 (1H), 7.94 (1H) ppm.

Example 33

(1S,3S(Z),7S,10R,11S,12S,16S)-7,11-Dihydroxy-3-(2-methyl-
benzoxazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (A) and
(1R,3S(Z),7S,10R,11S,12S,16R)-7,11-dihydroxy-3-(2-methyl-
benzoxazol-5-yl)-10-(prop-2-en-1-yl)-8,8,12,16-tetramethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione (B)

Analogously to Example 10, 20 mg (39 μ mol) of the compound that is produced according to Example 21 is reacted, and after working-up and purification, 11.2 mg (21 μ mol, 54%) of title compound A and 2.9 mg (5.5 μ mol, 14%) of title compound B are isolated in each case as a colorless oil.

$^1\text{H-NMR}$ (CDCl_3) of A: δ = 0.98 (3H), 1.02 (3H), 1.19-1.78 (7H), 1.22 (3H), 1.30 (3H), 2.15 (2H), 2.28 (1H), 2.33-2.60 (4H), 2.64 (3H), 2.92 (1H), 3.58 (1H), 3.69 (1H), 4.18 (1H), 4.29 (1H), 5.01 (1H), 5.08 (1H), 5.72 (1H), 6.14 (1H), 7.31 (1H), 7.47 (1H), 7.64 (1H) ppm.

Example 34

In Vitro Activity of Epothilone Derivatives on Human Tumor Cell Lines

a) IC_{50} values [nM] for the growth inhibition of human MCF-7-breast and multi-drug-resistant NCI/ADR-carcinoma cell lines of

5

10

epothilone derivatives with 13Z-unsaturated in the crystal-violet
assay in comparison to Taxol.

15

20

25

30

35

40

45

50

55

Table 1:

Compound	MCF-7	NCl/ADR	Selectivity*
Taxol	3.5	> 100	> 28.6
Example 1	30	75	2.5
Example 2	25	70	2.8
Example 9	17	41	2.4
Example 13	34	n.d.	
Example 15	25	n.d.	
Example 21	32	n.d.	
Example 23	11	62	5.6
Example 27	25	41	1.6

* Selectivity = $IC_{50} - (NCl/ADR) : IC_{50} (MCF-7)$; n.d.: not yet determined

The compounds of applicants' invention have a significantly higher active strength in comparison to Taxol. All the compounds according to the invention which were tested show an action on the multi-drug-resistant cell line NCl/ADR not exhibited by Taxol.

b) IC_{50} values [nM] for the growth inhibition of human MCF-7-breast- and multi-drug-resistant NCl/ADR carcinoma cell lines of epothilone derivative with 13,14- α -epoxide, which was formed from the 13-Z-configured double bond in the crystal-violet assay in comparison to Taxol.

Table 2:

Compound	MCF-7	NCl/ADR	Selectivity*
Taxol	3.5	> 100	> 29
Example 4A	1.3	9.1	7.0
Example 5A	3.1	3.8	1.2
Example 10A	1.2	3.6	3.0
Example 25	2.3	11	4.8

*Selectivity = $IC_{50} - (NCl/ADR) : IC_{50} - (MCF-7)$

In contrast to Taxol, all compounds of the invention show an action on the multi-drug-resistant cell line NCl/ADR.

c) IC_{50} values [nM] for the growth inhibition of human MCF-7-breast- and multi-drug-resistant NCl/ADR-carcinoma cell lines of epothilone derivatives with 13,14-epoxide, which was formed from the 13-E-configured double bond, in a crystal-violet assay in comparison to Taxol.

Table 3:

Compound	MCF-7	NC1/ADR	Selectivity*
Taxol	3.5	> 100	> 29
Example 6A or B	4.3	68	15.8
Example 12A or B	40	61	1.5
Example 12B or A	4.8	53	11.0
Example 26A or B	18	60	3.3

* Selectivity = $IC_{50} - (NC1/ADR) : IC_{50} - (MCF-7)$

In contrast to Taxol, all compounds show an action on the multi-drug-resistant cell line NC1/ADR.

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and, without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

Claims

5

10

15

20

25

30

35

40


45

50

55

Claims

1. An epothilone compound of formula I,



I,


in which

R^{1a} , R^{1b} are the same or different and mean hydrogen; C_1 - C_{10} alkyl, C_6 - C_{12} aryl, or C_7 - C_{20} aralkyl, each optionally substituted; or taken together a $-(CH_2)_m-$ group with m

$= 1, 2, 3, 4$ or 5 or a $-(CH_2)-O-(CH_2)-$ group;

R^{2a} means hydrogen, C_1 - C_{10} alkyl, C_6 - C_{12} aryl or C_7 - C_{20} aralkyl, each optionally substituted;

$-(CH_2)_{ra}-C\equiv C-(CH_2)_{pa}-R^{26a}$; $-(CH_2)_{ra}-C=C-(CH_2)_{pa}-R^{26a}$,



10



20

25

30

35

40

45

50

5

10

15

20

25

30

35

40

45

50

55

s stands for 1, 2, 3 or 4,

T stands for OR^{22} or Hal,

R^{22} stands for hydrogen or a protective group
 PG^4 ,

R^6 , R^7 each mean a hydrogen atom, or taken together an
additional bond or an oxygen atom,

G means a group $\text{X}=\text{CR}^8$ - or a bi- or tricyclic aryl
radical,

R^8 means hydrogen; halogen; CN; or a C_1 - C_{20} alkyl, C_6 - C_{12}
aryl, C_7 - C_{20} aralkyl group each of which can be
substituted,

X means an oxygen atom, two alkoxy groups OR^{23} , a C_2 - C_{10} -
alkylene- α,ω -dioxy group which can be straight-chain or
branched, H/OR^9 or a grouping $\text{CR}^{10}\text{R}^{11}$,

whereby

R^{23} stands for a C_1 - C_{20} alkyl radical,

R^9 stands for hydrogen or a protective group PG^x ,

R^{10} , R^{11} are the same or different and stand for

hydrogen; a C_1 - C_{20} alkyl, C_6 - C_{12} aryl or C_7 - C_{20}

aralkyl radical each optionally substituted; or

R^{10} and R^{11} together with the methylene carbon atom

jointly stand for a 5- to 7-membered carbocyclic

ring,

5

D-E means a group $-\text{CH}_2-\text{CH}_2-$, $-\text{O}-\text{CH}_2-$,

10

A-Y means a group $\text{O}-\text{C}(=\text{O})$, $\text{O}-\text{CH}_2$, $\text{CH}_2\text{C}(=\text{O})$, $\text{NR}^{29}-\text{C}(=\text{O})$, or
 $\text{NR}^{29}-\text{SO}_2$,

15

R^{29} means hydrogen or C_1-C_{10} alkyl,

Z means an oxygen atom or H/OR^{12} ,

whereby

20

R^{12} is hydrogen or a protective group PG^* ,

Hal means halogen.

25

2. A compound of formula I according to claim 1, wherein
 R^{2a} stands for a hydrogen atom.

30

3. A compound of formula I according to claim 1, wherein
 R^{1a} and R^{1b} in each case stand for a methyl group or together form
a dimethylene or trimethylene group.

35

40

4. A compound of formula I according to claim 1, wherein G
is a group $\text{X} = \text{CR}^8-$ where R^8 is a halogen atom or a nitrile
group.

45

5. A compound according to claim 4, wherein R^8 is a
fluorine atom.

50

55

5

10

6. A compound of formula I according to claim 1, selected from the group consisting of:

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-Dihydroxy-16- (1-methyl-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R), 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-chloro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopentyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopentyl-1-ethyl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-

30

8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopentyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopentyl-1-ethyl)-3-(2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopentyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione

50

55

5

10

15

20

25

30

35

40

45

50

55

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-Dihydroxy-16- (1-chloro-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-Dihydroxy-10- (prop-2-in-1-yl) -3- (1-chloro-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7-

40

(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -

30

8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

25

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, (10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-fluoro-2- (2-methylthiazol-4-

50

55

5

10

15

20

25

30

35

40

45

50

55

yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(2-oxacyclopropyl-1-ethyl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(2-oxacyclopropyl-1-ethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

20

25

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]

45

heptadecane-5,9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-

35

tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-methyl-2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

30

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (2- (2-methyloxazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-chloro-2- (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl)ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl)ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4,17-

40

dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-

50

55

5

10

trimethylene-12,16-dimethyl-4,17-dioxabicyclo[14.1.0]heptadecane-
5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-
methylthiazol-4-yl)ethenyl)-1-oxa-5,5-trimethylene-9,13-dimethyl-
7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-
in-1-yl)-3-(1-methyl-2-(2-methylthiazol-4-yl)ethenyl)-8,8-
trimethylene-12,16-dimethyl-4,17-dioxabicyclo[14.1.0]heptadecane-
5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-
methylthiazol-4-yl)ethenyl)-1-oxa-5,5-trimethylene-9,13-dimethyl-
7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-
in-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8-

35

trimethylene-12,16-dimethyl-4,17-dioxabicyclo[14.1.0]heptadecane-
5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-
methylthiazol-4-yl)ethenyl)-1-oxa-5,5-trimethylene-9,13-dimethyl-
7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-
en-1-yl)-3-(2-(2-methylthiazol-4-yl)ethenyl)-8,8-trimethylene-
12,16-dimethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

50

55

5

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2-(2-methylthiazol-4-yl)ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl) -3- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-

20

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (2- (2-methyloxazol-4-yl)ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl)ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl)ethenyl) -1-oxa-5, 5-trimethylene-9, 13-dimethyl-

50

7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

45

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-oxa-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

50

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-chloro-2- (2-methyloxazol-4-

55

5

10

yl)ethenyl)-8,8-trimethylene-12,16-dimethyl-4,17-
dioxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-
benzoxazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-
cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-
en-1-yl)-3-(1-methyl-2-(2-methyl-benzoxazol-5-yl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-
benzoxazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-
cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-
in-1-yl)-3-(1-methyl-2-(2-methyl-benzoxazol-5-yl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-
benzoxazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-
cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-
en-1-yl)-3-(1-methyl-2-(2-methyl-benzoxazol-5-yl)-8,8,12,16-
tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

45

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-
benzoxazol-5-yl)-1-oxa-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-
cyclohexadec-13-ene-2,6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-

35

4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (quinolin-2-yl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (quinolin-2-yl) - 8, 8, 12, 16-tetramethyl-

45

4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (quinolin-2-yl) -
1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-
methyl-but-2-en-1-yl) -3- (1-methyl-2- (quinolin-2-yl) -8, 8, 12, 16-
tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-
benzoxazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
en-1-yl) -3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) -8, 8, 12, 16-
tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-
benzoxazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -
cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-
in-1-yl) -3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) -8, 8, 12, 16-
tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2-methyl-
benzoxazol-5-yl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -
cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzoxazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzoxazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2-methyl-benzothiazol-5-yl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyl-benzothiazol-5-yl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-methyl-2-(2-methyl-benzothiazol-5-yl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-methyl-2-(2-methyl-benzothiazol-5-yl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-methyl-benzothiazol-5-yl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-methyl-benzothiazol-5-yl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-methyl-2-(2-methyl-benzothiazol-5-yl)-

50

55

5

10

8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(quinolin-2-yl))-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-in-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-in-1-yl)-3-(1-methyl-2-(quinolin-2-yl))-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-en-1-yl)-cyclohexadec-13-ene-2,6-dione

40

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-en-1-yl)-3-(1-methyl-2-(quinolin-2-yl))-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

45

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(quinolin-2-yl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione

50

55

5

252

10

NOT FURNISHED UPON FILING

15

20

25

30

35

40

45

50

55

5

253

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

254

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

255

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (2- (2-pyridyl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-fluoro-2- (2-pyridyl)ethenyl) -

30

8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-chloro-2- (2-pyridyl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-

45

dione

50

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

55

5

256

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

257

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-methyl-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.10]heptadecane-5,9-dione

20

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

25

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(prop-2-en-1-yl)-3-(1-fluoro-2-(2-pyridyl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-pyridyl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(prop-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

50

55

5

258

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl-3- (2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

259

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2-(2-pyridyl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10-(oxacyclopropylmethyl) -3- (1-chloro-2- (2-pyridyl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

260

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-in-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

261

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

262

10

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) -3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

263

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

264

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-methylthiazol-4-yl)ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl)ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

25

oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-methylthiazol-4-

50

55

5

10

15

20

25

30

35

40

45

50

55

yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R(RS),8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(oxacyclopropylmethyl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R(RS),11R,12S,16R/S)-7,11-dihydroxy-10-(oxacyclopropylmethyl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione
(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(but-3-in-1-yl)-cyclohexadec-13-ene-2,6-dione
(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(but-3-in-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

5

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-in-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (but-3-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

268

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (but-3-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (but-3-en-1-yl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopropyl-1-ethyl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (2-oxacyclopropyl-1-ethyl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

269

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (2-oxacyclopentyl-1-ethyl) -cyclohexadec-13-ene-2, 6-dione

35

40

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (2-oxacyclopentyl-1-ethyl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

270

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E) , 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

271

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

20

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

50

55

5

272

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

45

50

55

5

. 273

10

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl) - 3 - (1-fluoro-2 - (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

15

oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-chloro-2 - (2-methyloxazol-4-yl)ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7-(oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10-(oxacyclopropylmethyl) - 3 - (1-chloro-2 - (2-methyloxazol-4-yl)ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-methyl-2 - (2-pyridyl)ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7 - (but-3-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10 - (but-3-in-1-yl) - 3 - (1-methyl-2 - (2-pyridyl)ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16 - (1-fluoro-2 - (2-pyridyl)ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7 - (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10 - (prop-2-in-1-yl) - 3 - (1-fluoro-2 - (2-pyridyl)ethenyl) - 8, 8-trimethylene-

50

12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

55

5

274

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R(RS) 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

15

oxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) - 1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (prop-2-in-1-yl) - 3- (2- (2-methyloxazol-4-yl) ethenyl) - 8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

276

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-in-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-in-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

20

oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-methyl-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (prop-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (prop-2-en-1-yl) -3- (1-fluoro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12-16-dimethyl-4-aza-17-

45

oxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

277

10

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-

20

oxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R(RS), 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-aza-5, 5-trimethylene-9, 13-dimethyl-7- (oxacyclopropylmethyl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R(RS), 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (oxacyclopropylmethyl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8-trimethylene-12, 16-dimethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -

45

8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

50

55

5

278

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-fluoro-2- (2-pyridyl) ethenyl) -

30

8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-chloro-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-dioxobicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

10

15

20

25

30

35

40

45

50

55

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-fluoro-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4, 17-

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-methylthiazol-4-yl) ethenyl) - 1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-chloro-2- (2-methylthiazol-4-

5

280

10

yl)ethenyl)-8,8,12,16-tetramethyl-4,17-

dioxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-

15

methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-

methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-

20

methyl-but-2-en-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-

yl)ethenyl)-8,8,12,16-tetramethyl-4,17-

dioxabicyclo[14.1.0]heptadecane-5,9-dione

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-

methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-

30

methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-

methyl-but-2-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-

35

8,8,12,16-tetramethyl-4,17-dioxabicyclo[14.1.0]heptadecane-5,9-dione

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-

40

methyloxazol-4-yl)ethenyl)-1-oxa-5,5,9,13-tetramethyl-7-(3-

methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-

45

methyl-but-2-en-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-

yl)ethenyl)-8,8,12,16-tetramethyl-4,17-

50

dioxabicyclo[14.1.0]heptadecane-5,9-dione

55

5

281

10

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -1-oxa-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

15

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-chloro-2- (2-methyloxazol-4-yl) ethenyl) -8, 8, 12, 16-tetramethyl-4, 17-

20

dioxabicyclo[14.1.0]heptadecane-5, 9-dione

25

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-methyl-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

30

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (1-methyl-2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

35

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

40

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) -7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) -3- (2- (2-pyridyl) ethenyl) -8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

45

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) -4, 8-dihydroxy-16- (1-fluoro-2- (2-pyridyl) ethenyl) -1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) -cyclohexadec-13-ene-2, 6-dione

50

55

5

282

10

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-fluoro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

15

20

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-chloro-2- (2-pyridyl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

25

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-chloro-2- (2-pyridyl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

30

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

35

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (1-methyl-2- (2-methylthiazol-4-yl) ethenyl) - 8, 8, 12, 16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5, 9-dione

40

(4S, 7R, 8S, 9S, 13E/Z, 16S(E)) - 4, 8-dihydroxy-16- (2- (2-methylthiazol-4-yl) ethenyl) - 1-aza-5, 5, 9, 13-tetramethyl-7- (3-methyl-but-2-en-1-yl) - cyclohexadec-13-ene-2, 6-dione

45

(1S/R, 3S(E), 7S, 10R, 11R, 12S, 16R/S) - 7, 11-dihydroxy-10- (3-methyl-but-2-en-1-yl) - 3- (2- (2-methylthiazol-4-yl) ethenyl) -

50

55

5

283

10

8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

15

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

20

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-fluoro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-

25

oxabicyclo[14.1.0]heptadecane-5,9-dione

30

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

35

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-chloro-2-(2-methylthiazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-

oxabicyclo[14.1.0]heptadecane-5,9-dione

40

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

45

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-methyl-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-

50

oxabicyclo[14.1.0]heptadecane-5,9-dione

55

5

284

10

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

15

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

20

25

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione

30

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-fluoro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

35

(4S,7R,8S,9S,13E/Z,16S(E))-4,8-dihydroxy-16-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-1-aza-5,5,9,13-tetramethyl-7-(3-methyl-but-2-en-1-yl)-cyclohexadec-13-ene-2,6-dione and

40

(1S/R,3S(E),7S,10R,11R,12S,16R/S)-7,11-dihydroxy-10-(3-methyl-but-2-en-1-yl)-3-(1-chloro-2-(2-methyloxazol-4-yl)ethenyl)-8,8,12,16-tetramethyl-4-aza-17-oxabicyclo[14.1.0]heptadecane-5,9-dione

45

50

55

5

285

10

7. A compound of the formula I of claim 1, wherein Z is oxygen.

15

8. A compound of the formula I of claim 1, wherein R^{1a} and R^{1b} taken together form a trimethylene group.

20

9. A compound of the formula I of claim 1, wherein R^{2a} is hydrogen.

25

10. A compound of the formula I of claim 1, wherein R^{2a} is hydrogen and R^{2b} is allyl, prop-2-ynyl, but-3-ynyl or but-3-enyl.

30

11. A compound of the formula I of claim 1, wherein R^{3a} and R^{3b} taken together are H/OH.

35

12. A compound of the formula I of claim 1, wherein R⁴ is methyl, ethyl, propyl, n-butyl, i-butyl, t-butyl or benzyl.

40

13. A compound of the formula I of claim 1, wherein R⁵ is H, C₁₋₅ alkyl or -(CH₂)_s-T where T is OH, F or Cl and s is 1 or 2.

45

50

55

5

286

10

14. A compound of the formula I of claim 1, wherein G is a bicyclic heteroaryl radical with at least one nitrogen atom in the bicyclic ring.

15

15. A compound of claim 14, wherein G is a 2-methylbenzoxazol-5-yl or 2-methylbenzothiazol-5-yl radical.

20

25

16. A compound of the formula I of claim 1, wherein G is a group $X=CR^8-$, where R^8 is C_1-C_4 alkyl, hydrogen, F, Cl or Br; and X is $CR^{10}R^{11}$ where one of R^{10} and R^{11} is hydrogen and the other is a heterocyclic aryl radical.

30

17. The compound of claim 16, wherein the heterocyclic aryl radical is 2-methylthiazol-4-yl or 2-pyridyl.

35

18. A compound of the formula I of claim 1, wherein -A-Y- is $-O-C(=O)-$ or $-NR^{29}-C(=O)-$ where R^{29} is hydrogen or C_1-C_3 alkyl.

40

19. A pharmaceutical composition comprising a compound of the formula I of claim 1 and at least one pharmaceutically acceptable adjuvant or vehicle.

45

50

55

20. A composition according to claim 19 in the form of a dosage unit containing 0.1-100 mg of the compound of the formula I of claim 1.


21. A method for treating a disease or condition associated with cell growth, division and/or proliferation which comprises administering a compound of the formula I of claim 1.

22. The method of claim 21, wherein the compound of formula I is administered to a human in a dosage of 0.1-100 mg/day.

23. A prodrug for a compound of the formula I of claim 1 which comprises an entity which provides, in vivo, a compound of the formula I of claim 1.

24. A method for preparing a compound of the formula I of claim 1, which comprises:

cyclizing a compound of the formula ABC-1 or ABC-2



5

288

10

wherein R^{14} , Z, R^{3a} , R^4 , D, E, R^5 , R^6 , R^7 and G have the meaning given in claim 1,

15

$R^{1a'}$, $R^{1b'}$, $R^{2a'}$ and $R^{2b'}$ have the meanings given for R^{1a} , R^{1b} , R^{2a} and R^{2b} in claim 1

R^{13} means CH_2OR^{13a} , CH_2-Hal , CHO , CO_2R^{13b} or $COHal$,

20

R^{13a} means hydrogen, SO_2 -alkyl, SO_2 -aryl, SO_2 -aralkyl or together a $-(CH_2)_q$ - group or together a $CR^{15a}R^{15b}$ group,

R^{13b} means hydrogen, C_1 - C_{20} alkyl, C_6 - C_{12} aryl, C_7 - C_{20} aralkyl, each optionally substituted,

25

R^{15a} , R^{15b} are the same or different and mean hydrogen, C_1 - C_{10} alkyl, aryl, C_7 - C_{20} aralkyl or together a $-(CH_2)_q$ group,

30

$PG^{14'}$ is hydrogen or a protecting group,

R^{20} means halogen, N_3 , NHR^{29} , a hydroxy group, a protected hydroxy group $O-PG^2$, a protected amino group $NR^{29}PG^2$, a C_1 - C_{10} alkylsulfonyloxy group, which optionally can be perfluorinated, a benzyloxy group that is optionally substituted by C_1 - C_4 alkyl, nitro, chlorine or bromine, an $NR^{29}SO_2CH_3$ group, an $NR^{29}C(=O)CH_3$ group, a $CH_2-C(=O)-CH_3$ group,

40

R^{30} means hydrogen,

45

R^{31} means hydroxyl, or


50

55


R^{30} , R^{31} together mean an oxygen atom or a C_2 - C_{10} alkylene- α,ω -
 dioxy group, which can be straight-chain or branched,
 or

R^{30} , R^{31} independently mean a C_1 - C_{10} alkoxy group.


25. The method of claim 24, wherein the compound of ABC-1
 or ABC-2 is prepared by reacting an intermediate AB-1 or AB-2





with fragment C_f ,



or reacting intermediate BC



with fragment A-1 or A-2



or

A-1

A-2

wherein

V means an oxygen atom, two alkoxy groups OR^{17} , a C_2-C_{10} alkylene- α,ω -dioxy group, which can be straight-chain or branched or H/OR^{16} ,

W means an oxygen atom, two alkoxy groups OR^{19} , a C_2-C_{10} alkylene- α,ω -dioxy group, which can be straight-chain or branched or H/OR^{18} ,

G' means a group $X=CR^{8'}$ -, a bicyclic or tricyclic aryl radical,

$R^{8'}$ has the meaning already mentioned in general formula I for R^8 ,

$R^{7'}$ means a hydrogen atom,

R^{13a} means hydrogen, SO_2 -alkyl, SO_2 -aryl, SO_2 -aralkyl or together a $-(CH_2)_6-$ group or together a $CR^{15a}R^{15b}$ group,

R^{11b} means hydrogen, C_1-C_{20} alkyl, C_6-C_{12} aryl, C_7-C_{20} aralkyl, each optionally substituted,

5

10

15

20

25

30

35

40

45

50

55

R^{15a} , R^{15b} are the same or different and mean hydrogen, C_1 - C_{10} alkyl, aryl, C_7 - C_{20} aralkyl or together a $-(CH_2)_q$ group,

X means an oxygen atom, two alkoxy groups OR^{23} , a C_2 - C_{10} alkylene- α,ω -dioxy group, which can be straight-chain or branched, H/OR^9 or a grouping $CR^{10}R^{11}$, whereby

R^{23} stands for a C_1 - C_{20} alkyl radical,

R^9 stands for hydrogen or a protective group PG^1 ,

R^{10} , R^{11} are the same or different and stand for hydrogen, a C_1 - C_{20} alkyl, C_6 - C_{12} aryl, or C_7 - C_{20} aralkyl radical, each optionally substituted, or R^{10} and R^{11} together with the methylene carbon atom commonly stand for a 5- to 7-membered carbocyclic ring, and

R^{21} means a hydroxy group, halogen, a protected hydroxy group OPG^1 , a phosphonium halide radical $PPh_3^+Hal^-$ (Ph = phenyl; Hal = F, Cl, Br, I), a phosphonate radical $P(O)(OQ)_2$ (Q = C_1 - C_{10} alkyl or phenyl) or a phosphine oxide radical $P(O)Ph_2$ (Ph = phenyl)

$R^{14'}$ means hydrogen, OR^{14a} , Hal, OSO_2R^{14b} ,

R^{14a} means hydrogen, SO_2 -alkyl, SO_2 -aryl, SO_2 -aralkyl or together a $-(CH_2)_o$ - group or together a $CR^{15a}R^{15b}$ group,


R^{14b} means hydrogen, C_1 - C_{20} alkyl, C_6 - C_{12} aryl, C_7 - C_{20} aralkyl, each optionally substituted,

o means 2 to 4,

q means 3 to 6, and

the other variables are as defined above.

26. The method of claim 25, wherein the intermediates AB-1 and AB-2 are prepared by reacting fragments A-1, A-2 or C₁ with fragment B₁



B₁

wherein R^{3a'}, R^{4'} and R^{5'} have the meanings given for R^{3a}, R⁴ and R⁵, respectively.

INTERNATIONAL SEARCH REPORT

Inter national Application No
PCT/IB 00/00657

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D493/04 C07D405/06 C07D417/06 C07D405/14 C07D413/06
A61K31/335

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 07692 A (SCHERING) 18 February 1999 (1999-02-18) cited in the application claims; examples 24,27,34 ---	1-26
Y	WO 98 38192 A (GESELLSCHAFT FÜR BIOTECHNOLOGISCHE FORSCHUNG) 3 September 1998 (1998-09-03) cited in the application claims; examples --- -/--	1-26

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

27 September 2000

Date of mailing of the international search report

23. 10. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/IB 00/00657

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	K. NICOLAOU ET. AL.: "Gezielt entworfene Epothilone: kombinatorische Synthese, Induktion der Tubulin-Polymerisation und cytotoxische Wirkung gegen taxolresistente Tumorzellen. " ANGEWANDTE CHEMIE, vol. 109, no. 19, 2 October 1997 (1997-10-02), pages 2181-7, XP002148547 Weinheim, DE cited in the application tables 1,2	1-26
A	K. NICOLAOU ET. AL. : "Synthesis of Epothilones A and B in solid and solution phase." NATURE, vol. 387, 8 May 1997 (1997-05-08), pages 268-72, XP002148548 cited in the application whole article	1-26

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 00/00657

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 21 and 22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/IB 00/00657

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9907692 A	18-02-1999	DE 19735574 A	11-02-1999
		DE 19735575 A	11-02-1999
		DE 19735578 A	11-02-1999
		DE 19748928 A	29-04-1999
		DE 19749717 A	06-05-1999
		DE 19751200 A	20-05-1999
		DE 19813821 A	23-09-1999
		AU 9340998 A	01-03-1999
		EP 1005465 A	07-06-2000
WO 9838192 A	03-09-1998	AU 6724998 A	18-09-1998
		BR 9807742 A	22-02-2000
		CN 1248974 T	29-03-2000
		DE 19880193 D	24-08-2000
		EP 0975638 A	02-02-2000
		NO 994071 A	21-10-1999
		PL 335329 A	25-04-2000
		ZA 9801575 A	08-09-1998